



Incorporating mortality into habitat selection to identify secure and risky habitats for savannah elephants



C.L. Roever^{a,*}, R.J. van Aarde^{a,*}, M.J. Chase^{a,b,c}

^a Conservation Ecology Research Unit, Department of Zoology and Entomology, University of Pretoria, Hatfield 0083, South Africa

^b Elephants Without Borders, PO Box 682, Kasane, Botswana

^c San Diego Zoo Institute for Conservation Research, San Diego Zoo Global, 15600 San Pasqual Valley Road, Escondido, CA 92027, United States

ARTICLE INFO

Article history:

Received 8 November 2012

Received in revised form 30 January 2013

Accepted 8 April 2013

Keywords:

African savannah elephant

Anthropogenic features

Habitat quality

Habitat selection

Mortality

Source and sink

ABSTRACT

Empirical models of habitat selection are increasingly used to guide and inform habitat-based management plans for wildlife species. However, habitat selection does not necessarily equate to habitat quality particularly if selection is maladaptive, so incorporating measures of fitness into estimations of occurrence is necessary to increase model robustness. Here, we incorporated spatially explicit mortality events with the habitat selection of elephants to predict secure and risky habitats in northern Botswana. Following a two-step approach, we first predict the relative probability of use and the relative probability of mortality based on landscape features using logistic regression models. Combining these two indices, we then identified low mortality and high use (primary habitat) and areas of high mortality and high use (primary risk). We found that mortalities of adult elephants were closely associated with anthropogenic features, with 80% of mortalities occurring within 25 km of people. Conversely, elephant habitat selection was highest at distances of 30–50 km from people. Primary habitat for elephants occurred in the central portion of the study area and within the Okavango Delta; whereas risky areas occurred along the periphery near humans. The protected designation of an area had less influence on the proportion of prime habitat therein than did the locations of the area in relation to human development. Elephant management in southern Africa is moving towards a more self-sustaining, habitat-based approach, and information on selection and mortality could serve as a baseline to help identify demographic sources and sinks to stabilize elephant demography.

© 2013 Elsevier Ltd. All rights reserved.

1. Introduction

Using telemetry data to predict habitat utilization is a common practice in ecology, and the results are often used as a guide to protect and preserve habitats important to wildlife species (e.g. Johnson et al., 2004; Meyer et al., 1998). However, habitat utilization is only one aspect within a complex set of factors that ultimately relate to individual fitness. Selection, for example, could be maladaptive, whereby individuals select areas that ultimately increase their mortality risk or decrease their reproductive success (Battin, 2004; Delibes et al., 2001). To avoid the misclassification of highly used habitats as high-quality habitats, it is necessary to temper estimations of habitat selection with some indication of fitness or risk, particularly when habitat selection appraisals are to be used as an impetus for conservation action.

The main limitation of incorporating indices of fitness into animal occurrence models is that spatially-explicit fitness data is often difficult to obtain (Nielsen et al., 2006). Studies that

incorporate offspring survival are most common for avian species, where fledging success at the nest site can readily be established (Aldridge and Boyce, 2007; Donovan and Thompson, 2001). However, with mammalian species the estimation of recruitment and survival are generally less straightforward. Changes in the demographic parameters of large mammals in particular occur over relatively long timescales because of their multi-year or multi-decade life spans and their low reproductive and mortality rates. Most large mammals are also highly mobile with large ranging patterns, making it difficult to relocate individuals to monitor survival and fecundity. Studies linking reproductive success to habitats have been most successful in ungulates on islands and other closed systems (McLoughlin et al., 2007, 2008). Given limitations in time, resources, study species, and study area, researchers resort to data that are more readily available, such as mortality location data (Dzialak et al., 2011; Falcucci et al., 2009; Nielsen et al., 2006).

Just as live animals inform habitat selection estimations, carcass locations provide spatially explicit information on where animals die and can be used to inform the riskiness of habitats (Nielsen et al., 2006). However, mortality location data is often underutilized in the literature. African elephants, for example, have

* Corresponding author. Tel.: +27 12 420 2753; fax: +27 12 420 4523.

E-mail addresses: mail@carrieroever.com (C.L. Roever), rjvaarde@zoology.up.ac.za (R.J. van Aarde).

carcasses that are easily visible from the air for several years after death (Douglas-Hamilton and Hillman, 1981), and while carcasses have been used to inform local mortality rates (e.g. Dudley et al., 2001; Dunham, 2008) and CITES status (e.g. Wittemyer et al., 2013), we found no peer-reviewed study exploiting the spatial location of carcasses. Knowing where animals die can provide valuable insight into risky landscapes, which is helpful information to guide conservation and management plans. Grizzly bears mortalities in Alberta, for example, were concentrated around roads or hiking trails (Benn and Herrero, 2002), prompting calls to regulate human access in grizzly bear habitats (Alberta Sustainable Resource Development and Alberta Conservation Association, 2010). Relating mortality locations to habitat selection models has also been used to better inform habitat-based management plans (i.e. Nielsen et al., 2006).

Habitat heterogeneity contributes to the spatial pattern of use and mortalities of elephants, and describing those patterns is particularly important in Africa, where the management of savannah elephants is a continuing concern (van Aarde and Jackson, 2007). Hunting and poaching in the late 19th and early to mid 20th century reduced some populations to near extinction (Roth and Douglas-Hamilton, 1991; Whyte et al., 2003), but actions taken to decrease poaching in the mid to late-20th century were largely successful in southern Africa (Whyte et al., 2003). The decline and subsequent recovery of elephant populations also may explain changes in woodland habitat (Guldmond and van Aarde, 2008; Nasseri et al., 2011), prompting concern for elephant-related tree damage (van Aarde et al., 2008). Their role as ecosystem engineers and the susceptibility to population decline from legal and illegal hunting makes it particularly important to quantify how habitats contribute to elephant habitat selection and mortality risk.

Using the information theoretic approach, we modelled habitat selection from elephant occurrence data obtained during aerial surveys. We then created an index of habitat use by elephants. Following a similar procedure, we next modelled the relative probability of elephant mortality using the locations of elephant carcasses. Combining the relative probability of use and mortality indexes, we then defined areas of high use and low mortality as secure habitats, and areas of high use and high mortality as risky areas. By interrelating conditions where elephants live with where they die, we can begin to establish a habitat-based approach to elephant management and work towards understanding and maintaining natural regulatory processes where needed, as proposed by van Aarde and Jackson (2007) and supported by others (Chamaillé-Jammes et al., 2008). These indices, while not a direct measure of demographic sources and sinks, do provide insights for the prioritization of conservation actions and can serve as a baseline to direct future studies into elephant demography.

2. Methods

2.1. Study area

The study area in northern Botswana incorporated an area of 74,355 km². Study area boundaries to the north and east coincide with national borders for Namibia, Zambia, and Zimbabwe, respectively (Fig. 1). Twenty percent of the study area was protected within the confines of Chobe National Park (NP), Makgadikgadi NP, Moremi Game Reserve (GR), and Nxai Pan NP, while an additional 65% occurred within Wildlife Management Areas (WMAs; Chase, 2011). WMAs were designated with a two letter code and number and, for the purpose of this study, were grouped into study




Fig. 1. Map of the study area located in northern Botswana. The area included three national parks, one game reserve, and multiple wildlife management areas which were grouped and referred to as study regions.

Table 1

Structure of the AIC candidate models evaluating elephant habitat use (elephant location vs. random location), elephant mortality risk (mortality location vs. random location), and difference in selection (mortality location vs. elephant location). Distance to water and distance to road were correlated; therefore, they could not be used in the same model.

Model name	Model
1. Null	
2. Water	Distance to water ^a
3. Nutrients	Distance to water ^a + Proportion tree ^a
4. Water, food, and slope	Slope + Distance to water ^a + Proportion tree ^a
5. Human presence	Distance to humans ^{b,c} + Distance to road ^b
6. Cover and humans	Proportion tree ^a + Distance to humans ^{b,c} + Distance to road ^b
7. Global model with water	Slope + Distance to water ^a + Proportion tree ^a + Distance to humans ^{b,c}
8. Global model with roads	Slope + Proportion tree ^a + Distance to humans ^{b,c} + Distance to road ^b

^a Quadratic in all analysis.

^b Quadratic in the elephant vs. random analysis.

^c Quadratic in the mortality vs. elephant analysis.




Fig. 2. Five habitat states categorized based on relative probability of use (10 ordinal bins from 1-low to 10-high) and relative probability of mortality (10 ordinal bins from 1-low to 10-high) for elephants. This figure was adapted from Nielsen et al. (2006).

regions (Fig. 1). Legal trophy hunting of elephant males was permitted in WMAs between the months of April and September, with a quota of 400 males in 2010 and 306 taken. Most of the terrain within the study area was flat, with the steepest slopes of eight degrees occurring along the Chobe River. The vegetation consisted primarily of deciduous dry woodlands and interspersed grasslands (Gaughan et al., 2012). Kasane, in the northeastern corner of the study area, was the largest town, and outside of this town, human settlements occurred in small villages along roadways mainly on the periphery of the study area. Permanent human settlements were prohibited in national parks; however lodges, campsites,

and park offices were located within park boundaries. Roads also occurred throughout at a density of 0.073 km/km².

2.2. Elephant data

Aerial surveys to locate elephant carcasses and live elephants were conducted from June to December 2010. Aerial survey methodology followed procedures established by Norton-Griffiths (1978), whereby parallel strip-transects were systematically flown in a fixed-wing plane and animal locations were recorded. Transects were flown at a speed of 160 km/h and 300 feet above ground level. The study area was divided into 42 sampling units, and transects within each sampling unit were surveyed once over a period of 1 day to reduce duplicate counting of animals. The distance between strip-transects varied by sampling unit to minimize sampling effort (for detailed methods see Chase, 2011). Sampling units expected to have high to moderate elephant density were surveyed more intensively (2–4 km, $n = 38$) than those with low elephant densities (8 km, $n = 3$). However, NG26 was surveyed at a 1 km density at the request of local stake holders. The width of strip transects extended approximately 400 m (two observers covering 200 m); consequently, survey intensities of 1, 2, 4, and 8 km accounted for 40%, 20%, 10%, and 5% coverage of the sampling unit, respectively. The differing sampling intensities were not biased towards a particular habitat type or landscape feature and would, therefore, not bias habitat selection models (see Appendix A in Supplementary material). Orientation and spacing of flight paths were generated using DNR Garmin Sampling Extension in ArcView 3.2 (ESRI, Redlands, California).

The timing of aerial surveys coincided with the peak dry season to increase visibility through the tree canopy. The location, time, sex, and number of elephants were recorded. The sex and age

Table 2

Beta coefficients (β) and standard errors (SE) for the top-ranked AIC models. An asterisk (*) was used to indicate where the confidence intervals did not overlap with zero. Model fit using k -fold cross validation and the Spearman rank correlation coefficient (r_s) is also presented.

	Elephant vs. random		Mortality vs. random		Mortality vs. elephant	
	β	SE	β	SE	β	SE
Distance to water ^a	0.73	0.55	5.31	1.80*	4.29	1.90*
(Distance to water ^a) ²	-0.15	0.02*	-0.28	0.06*	-0.12	0.06
Slope	0.15	0.04*	0.40	0.07*	0.27	0.09*
Proportion tree	6.24	0.67*	13.36	2.23*	4.38	2.48
(Proportion tree) ²	-17.98	2.28*	-38.83	8.01*	-14.86	8.34
Distance to humans ^a	2.92	0.46*	-4.52	0.46*	-8.05	1.45*
(Distance to humans ^a) ²	-0.04	0.01*	-	-	0.05	0.03
Model fit (r_s)	1.00		0.96		0.87	

^a Coefficient value and standard error multiplied by 100 for ease of interpretation.

composition of individuals in groups were used to distinguish breeding herds from male herds which contained no females or juveniles. For elephant remains, the age of the carcass was categorized as either a recent carcass (i.e. flesh still present and the ground still moist indicating that the animal died less than a year ago) or an old carcass (i.e. clean bones, dry ground, and vegetation reestablishing in the rot patch) (Douglas-Hamilton and Hillman, 1981). From the air, observers also recorded whether multiple carcasses were observed in the same location to indicate illegal hunting. While observers also recorded whether tusks had been chopped or removed, park officials also removed tusks from any carcass they find, as did legal hunters. However, natural mortality

and hunter related deaths were generally isolated events; therefore, clustering of carcasses provided a better indication of poaching than did presence of tusks. Indications of illegal hunting were only noted in 18 instances, so low sample size precluded further analysis on poaching.

2.3. Modelling animal locations

We used a resource selection function (RSF) model to identify habitats associated with live elephant and elephant mortality locations. A RSF model compares animal locations to random locations within a logistic regression framework, using landscape




Fig. 3. Relative probability of use (A) and relative probability of mortality and (B) for elephants based on logistic regression models and binned into 10 ordinal classes. Relative probability of mortality represents year-round mortality events, whereas relative probability of use pertains to dry season (June–December) occurrence. Black lines indicate jurisdictional boundaries.




Fig. 4. Relative probability of occurrence for live elephant (solid black) and elephant carcass (dashed gray) locations as a function of distance to water, slope, and distance to humans.

Table 3

Percent composition of habitat states occurring within each study region.

	Non-habitat	Secondary habitat	Secondary risk	Primary habitat	Primary risk	Total area (km ²)
Chobe NP	14.6	14.5	26.1	14.8	30.0	10,751
Makgadikgadi NP	66.4	20.1	10.6	1.0	2.0	5018
Moremi GR	13.6	18.1	7.1	51.1	10.2	4889
Nxai Pan NP	35.8	27.9	8.1	9.5	18.7	2518
CH – east	42.1	9.8	19.3	9.0	19.7	6977
CH – north	39.1	0.5	43.2	0.1	17.2	2884
CT	84.1	3.0	11.2	0.4	1.3	10,002
NG – north	42.6	14.7	18.7	15.3	8.7	19,683
NG – south	10.4	33.1	7.0	36.5	13.0	8375
Total area	42.0	14.6	16.2	15.0	12.3	74,354




Fig. 5. Habitat states for elephants in northern Botswana. Non-critical habitats represent areas of low use by elephants. Primary and secondary habitats represent areas with low mortality and high to moderate use, respectively. Primary and secondary risk areas represent regions with high mortality and high to moderate use, respectively.

factors as independent variables. We used a design I approach because individual animals were not uniquely identified and were only sampled once (Manly et al., 2002). A design I approach is made at the population level, where used resources units are sampled for the entire study area (Manly et al., 2002). We then developed three habitat-based models. First, using the live elephant observations, we estimated habitat selection by elephants. Each elephant herd accounted for one observation, resulting in 3040 live elephant observations. We compared these to random

points generated at a density of 1 point per 3 km² across the study area, for a total of 24,785 random locations. Next, we estimated the habitat-specific probability of elephant mortality using the 341 elephant carcass observations. The same set of random points was used for the mortality model as in the elephant habitat selection model. Finally, we compared elephant carcass locations to live elephant locations. This model accounts for the reality that elephants can only die in areas where they occur (Nielsen et al., 2004).

Covariates used in the estimation of the three habitat selection models pertained to water, slope, tree cover, human presence, and roads (based on Roever et al., 2012). Surface water was located using data from Tracks4Africa (2010) and was visually validated against Landsat imagery. Missing water bodies were manually digitized into the water layer. Distance to water (km) was then calculated for each location. While surface water can vary substantially across the seasons, the time of death for mortality events was unknown; therefore, we used all water bodies in our estimation (analysis using dry season water availability is in Appendix B in Supplementary material). Because elephant locomotion is limited by steep slopes (Wall et al., 2006), slope was calculated in degrees using a 90×90 m digital elevation model (Jarvis et al., 2006). Proportion of tree cover was estimated using MODIS Vegetation Continuous Fields (Hansen et al., 2006), which assessed the proportion of tree cover at a 500×500 m resolution. To quantify permanent human use across the landscape, we used Landscan (2008) human population data. Landscan data estimates human use over a 24 h period, and includes permanent settlements as well as roads, recreation areas, and other areas where people occur but not necessarily sleep (Landscan, 2008). Areas with >16 people/km² were considered relevant for elephant space use (Hoare and Du Toit, 1999); therefore, these areas were identified and distance to these high human-use areas was calculated. Finally, road infrastructure data (Tracks4Africa, 2010) was used to determine the distance (km) of locations from roads. As most roads in the study area were secondary dirt roads, road type was not further differentiated. All geospatial analysis was completed using the Spatial Analyst extension of ArcGIS 10.0 (ESRI, Redlands, California) and Geospatial Modelling Environment (Beyer, 2011).

To determine which habitat variables influenced elephant use and death locations, Akaike's information criterion (AIC) was used to choose a top-model among eight a priori candidate models (Table 1; Burnham and Anderson, 2002). Prior to model selection, all variables were tested for correlations and for non-linearity. Using Pearson's correlation coefficient, variables with an $r > 0.60$ were not included in the same model. Correlations occurred between distance to roads and distance to water (Pearson's $r = 0.632$). All continuous variables were examined for nonlinearities using histograms and by examining changes in model fit when including a quadratic term. Model fit of the top-ranked model was evaluated using k -fold cross validation ($k = 5$) and the Spearman rank correlation coefficient (Boyce et al., 2002). Analyses were conducted in R software (R Core Team, 2012).

Using the resultant models for where elephants lived and where elephants died, we predicted the relative probability of use and the relative probability of mortality across the study area. The relative probabilities were then classified into 10 ordinal categories using the quantile (equal-area) method in ArcGIS 10.0, with the lowest and highest probability equal to 1 and 10, respectively (Nielsen et al., 2006).

2.4. Identifying habitat states

We followed the methods of Nielsen et al. (2006) to define five habitat states based on elephant use and mortality. Habitat states included non-habitat (low use), secondary habitat (moderate use, low mortality), primary habitat (high use, low mortality), secondary risk areas (moderate use, high mortality), and primary risk areas (high use, high mortality) (Fig. 2). We used the elephant data to define the boundaries between habitat states, using an a priori defined 80% cutoff. Consequently, where 80% of the live elephants occurred was categorized as habitat (use classes 5–10, inclusive) and where 80% of mortalities occurred was classified as high risk

(mortality classes 7–10, inclusive). We then calculated the percentage of each state that occurred within each study region, to ascertain the proportion of effective habitats present within each.

3. Results

The top-ranked model for all three habitat selection analyses (elephant vs. random, mortality vs. random, and mortality vs. elephant) was model 7 (the global model with water; see Appendix C in Supplementary material for full model results). This model had strong support in the elephant vs. random (weight = 1.00) and mortality vs. random (weight = 1.00) analyses. Conversely, in the analysis comparing carcass locations to live elephant locations, the global model with water (Model 7) had a weight of 0.74 and the global model with roads (Model 8) was the second ranked model with a weight of 0.23. These two models were similar in form except that they interchanged the correlated variables, distance to water and distance to road. In all three analysis, the top model provided good fit to the data using k -fold cross validation tested using the Spearman rank correlation coefficient ($r_s > 0.86$; Table 2).

For the elephant use model, selection was associated with intermediate distance from water, steep slopes, moderate tree cover, and further from humans (Table 2). These features were most prevalent in Moremi GR in the western portion of the study area and the WMAs surrounding the Okavango Delta (Fig. 3A). High elephant use was also prevalent in the central portion of the study area in NG 15, NG 18, and western Chobe NP. Selection patterns of elephant mortalities were similar to live elephant habitat use for most variables. Elephant mortalities were associated with areas of intermediate distance from water, steep slopes, and intermediate tree cover; however the magnitude of these selection coefficients varied from the elephant use model (Table 2). Counter to the elephant use model, mortalities were more closely associated with humans. Across the study area, high mortality areas were located along the periphery of the study area, near areas with higher levels of human occupation (Fig. 3B).

Results comparing elephant carcass locations to elephant use locations reveal that water, slope, and distance to humans differed significantly between the two. Compared to where they lived, elephants were more likely to die in areas slightly further from water and on steeper slopes (Fig 4); however, the most striking difference between use and mortality locations was proximity to humans. Live elephants selected areas which were at intermediate distances from humans, whereas the highest relative probability of mortality occurred in areas near humans. Overall, the predictions of models comparing mortality locations to either elephant locations or random locations were similar (Table 2), indicating that mortality was not closely associated with patterns of habitat selection but were instead more closely associated with some other factor, such as human presence.

Based on our classification of habitat states, 15.0% of the study area comprised primary habitat for elephants, while 12.3% were primary risk areas (Table 3). Secondary habitat, which had low mortality and moderate use by elephants, comprised 14.6% of the study area, whereas secondary risk areas, with high mortality and moderate use, composed 16.2%. Overall, 42.0% was classified as non-habitat. The study area contained three national parks and one game reserve. Of these, Chobe NP contained the highest proportion of primary (30.0%) and secondary (26.1%) risk areas of any study region, and Makgadikgadi NP had the second highest proportion of non-habitat of any study region and had little (1.0%) primary habitat for elephants. Conversely, Moremi GR had the highest percentage of primary habitat of any study region

(51.1%). WMAs also provide substantial amounts of primary and secondary habitat for elephants. The WMAs labelled NG north and south comprise 28,058 km² of land and they consist of 5666 km² (20%) of primary habitat (Fig. 5).

4. Discussion

Our results indicate that elephant use and mortality locations were spatially separated. Elephant mortality locations were concentrated in areas close to human settlements, with 80% of all elephant mortalities occurring within 25 km of high human use areas, an area that accounts for 52% of the study area. Conversely, 50% of live elephants were observed in that same area, and live elephants selected areas of intermediate distance from people (Fig. 4). Elephant mortalities were, therefore, not proportional to elephant distribution. This is positive for elephant conservation as it shows that elephants were not exhibiting maladaptive selection, whereby individuals select areas that ultimately increase their mortality risk. However, it also suggests that humans contribute to elephant mortalities in the study area. Nielsen et al. (2004) found a similar pattern with grizzly bears, whereby human-induced mortalities reflected patterns of human use rather than of grizzly bear use. However, cause of death for elephants can be difficult to ascertain (Moss et al., 2011). Tusk removal is not a reliable sign of human-caused mortality because game wardens also remove tusks on any carcass observed to discourage the trade of ivory. In this study only 18 of the 341 elephant carcasses showed obvious signs of poaching, and most of these (17 of 18) were within 25 km of high human use areas.

The use of carcass location data is particularly informative when the death of the individual is directly related to the habitat component in which it was found. For example, with predators or human hunters, the carcass of the animal will generally remain in close proximity to the source of risk. However, when the animal succumbs to starvation or a disease, the behaviour of the individual may change in response to the weakened state. Elephants, for example, are dependent on surface water for survival in northern Botswana (Metcalf and Kepe, 2008), and it has been anecdotally noted, although not statistically verified, that a sick elephant may remain within close proximity to a water source (Haynes, 1988). Here, we found that elephant mortalities were closely associated with water, but at a similar rate as live elephant locations (Fig. 4). Yet, our study period was not a particularly stressful year; therefore, years of drought may produce very different patterns of mortality. In such instances of nutrition stress and disease, the carcass location may prove to be less informative at a fine spatial scale as examined here. Across a larger spatial scale, however, carcass locations may still provide valuable information on the ability of the region to support elephant populations at the home range or landscape scale.

Most of our carcass observations were of adult elephants, possibly an artifact of their increased visibility from an airplane. Consequently, the relative probability of mortality described in this study is biased towards adults. Other studies examining juvenile mortality among elephants found in 89% of cases the cause of death was natural (i.e. from drought, starvation, disease, predation, etc.; Moss et al., 2011). Once an elephant reaches adulthood, the likelihood of natural mortality declines until the animal reaches the age of 40 years, when age-related factors again increase the risk of natural mortality (Moss et al., 2011). Because our observed carcass locations were of adults, it stands to reason that mortality was heavily biased towards human-dominated landscapes. A further bias in the data could exist as a consequence of the Okavango Delta. No study to date has examined elephant carcass longevity in a marshy environment like the delta, where water could aid in decay and dispersal or obscure carcass visibility from the air.

However, if the mortality events observed in this study are largely human-induced, then the low rate of mortality could simply be a consequence of decreased human presence in the delta. Nonetheless, more research needs to be done to verify the cause of death in adult elephants to better distinguish landscape factors associated with natural vs. human-induced mortality events.

Our study area in northern Botswana included three national parks, one game reserve, and many wildlife management areas. Because national parks have greater restrictions on animal and land-use practices, it is often assumed that these areas will provide lower mortality and more secure habitat for animals than would areas with fewer restrictions. However, this is not always the case in Africa where high human densities near park boundaries (Wittemyer et al., 2008) may increase incidences of poaching (Metzger et al., 2010; Milner-Gulland and Bennett, 2003) and human-wildlife conflicts (Newmark et al., 1994). Our results support this, as we found that areas designated as national parks and game reserves did not decrease mortality for elephants. The highest proportions of primary risk areas (high use and high mortality) occurred in Chobe NP, where 27% of our observed mortalities occurred; whereas the highest concentrations of primary habitat occurred in Moremi GP and NG-south (Table 3). Consequently, the protected designation of an area had less influence on elephant mortality than did the location of the protected area in relation to human occupation; therefore, increased human development in the study area will likely result in a greater abundance of risky areas for elephants.

We opted to use aerial survey data rather than telemetry data to make direct comparisons between elephant abundance and elephant mortality locations. However, the disadvantage of aerial survey data was that data were restricted to the dry season when decreased foliage on trees increased elephant visibility. Therefore, our conclusions of habitat states relate to dry season elephant use. The dry season is the limiting season for this mega-herbivore (Illiuss, 2006; Trimble et al., 2009), so we consider the dry season as it best reflects limitations in habitat suitability and ultimately best relates to fitness. Habitat selection patterns are likely to vary in the core wet season (Young et al., 2009), when elephants shift their diet to consume more grasses (Cerling et al., 2006; Codron et al., 2011) and are less dependent on permanent water sources (Harris et al., 2008; Young et al., 2009). Based on our knowledge of elephant use patterns in the area from GPS collar data, we suspect that areas designated as “non-habitat” will be most influenced, particularly in NG-north. This area has a high proportion of grasses and was used readily by elephants in the wet season. We also expect that elephants will be more closely associated with people during the dry season, as human settlements are also concentrated around permanent water, possibly increasing human-wildlife interactions and conflict. For patterns of elephant mortalities, results are less likely to vary with season because the longevity of elephant carcasses means that mortality events could have occurred during any season.

Habitat selection models are often used in the management and conservation of wildlife; however, maladaptive selection by wildlife can result in the conservation of habitats that do little to contribute to the local persistence of the species. By including an indication of risk, we can begin to temper habitat selection estimations to better define habitat suitability. Here we used mortality location data because the large body size and slow decay rate of elephant carcasses makes this data relatively straightforward to obtain (Douglas-Hamilton and Hillman, 1981); however other indicators which directly relate to individual fitness have also been used, such as the habitat selection of a predator (Hebblewhite et al., 2005), and the availability of a limiting resource (Bleich et al., 2010; Nielsen et al., 2010). The next step is to determine if these habitat selection and mortality models relate to demographic

responses by the population to determine if these areas of risk also act as attractive sinks (Battin, 2004; Delibes et al., 2001).

This study is the first to examine habitat-associations of adult elephant mortalities and provide spatially explicit predictions of secure and risky habitats. Understanding the possible habitat variables which could contribute to mortality is the first step towards developing a habitat-based management plan. Balancing factors that contribute to elephant fitness and mortality to stabilize growth is a management option that could help alleviate concerns of elephant overpopulation (van Aarde and Jackson, 2007). Past elephant management entailed fencing populations into protected parks and supplementing water, which often resulted in localized overpopulation (van Aarde and Jackson, 2007). To reduce elephant numbers, managers then resorted to regular but controversial culls (van Aarde et al., 2006). Conversely, the elephant population in northern Botswana has largely escaped this more hands-on management approach, and now provides a model for a self-sustaining population. The areas of primary habitat in the Okavango Delta and central portion of the study area were isolated from human development and could provide a source population to supplement potential risky areas closer to human development on the periphery if our proposed habitat states result in demographic response.

Because elephants have such large roaming areas, habitat-based management requires vast tracks of land to accommodate the habitat heterogeneity for secure and risky areas. To put the area in perspective, the study area is roughly the same size as the combined area of New Hampshire, Massachusetts, and Vermont in the United States. Yet our study area is just a portion of the larger Kavango–Zambezi Transfrontier Conservation Area, which stretches across 320,000 km² and five countries, and is home to some 220,000 elephants (Blanc et al., 2007). Across Africa, initiatives to create corridors, link protected areas, and establish transfrontier conservation areas may increase the amount of land available to elephants, while also increasing the heterogeneity of secure and risky habitats. This habitat heterogeneity, along with other factors such as long distance migrations, are natural regulatory processes which may have helped to limit elephant populations in the past and contribute to regional population stability (Owen-Smith, 2004; van Aarde and Jackson, 2007).

Acknowledgements

We would like to thank Elephants Without Borders, the International Fund for Animal Welfare, and the University of Pretoria for research funding. The aerial survey was sanctioned and supported by the Botswana Department of Wildlife and National Parks, through a grant administered by the Conservation Trust Fund (CTF/2010/56). Additional funding was received from the Zoological Society of San Diego, Madeleine and Jerry Delman Cohen, Mr. Brett Warren, Nathan Jamieson Memorial Fund, Abu Camp, and Mutual and Federal Insurance Botswana. Anna Songhurst is acknowledged for her support of the aerial survey over the Okavango Panhandle, and for making data available to complement this analysis. We thank Mr. Mike Holding of Afriscreen Films and Felix Goshier for flying the aerial survey. We appreciate the assistance of our aerial observers, Ms. Kelly Landen, Anna Songhurst and Allen Bekker. GIS support was provided by Tracks4Africa.

Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.biocon.2013.04.006>.

References

- Alberta Sustainable Resource Development and Alberta Conservation Association, 2010. Status of the Grizzly Bear (*Ursus arctos*) in Alberta: update 2010. Edmonton.
- Aldridge, C.L., Boyce, M.S., 2007. Linking occurrence and fitness to persistence: habitat-based approach for endangered greater sage-grouse. *Ecol. Appl.* 17, 508–526.
- Battin, J., 2004. When good animals love bad habitats: ecological traps and the conservation of animal populations. *Conserv. Biol.* 18, 1482–1491.
- Benn, B., Herrero, S., 2002. Grizzly bear mortality and human access in Banff and Yoho National Parks, 1971–1998. *Ursus* 13, 213–221.
- Beyer, H.L., 2011. Geospatial Modelling Environment. Spatial Ecology, LLC. <<http://www.spatial ecology.com/gme/>>.
- Blanc, J.J., Barnes, R.F., Craig, G., Dublin, H., Thouless, C., Douglas-Hamilton, I., Hart, J., 2007. African Elephant Status Report 2007: An Update from the African Elephant Database. International Union for Conservation of Nature, Gland.
- Bleich, V.C., Marshal, J.P., Andrew, N.G., 2010. Habitat use by a desert ungulate: predicting effects of water availability on mountain sheep. *J. Arid Environ.* 74, 638–645.
- Boyce, M.S., Vernier, P.R., Nielsen, S.E., Schmiegelow, F.K.A., 2002. Evaluating resource selection functions. *Ecol. Model.* 157, 281–300.
- Burnham, K.P., Anderson, D.R., 2002. Model Selection and Multimodel Inference: A Practical Information-theoretic Approach, second ed. Springer-Verlag, New York.
- Cerling, T.E., Wittemyer, G., Rasmussen, H.B., Vollrath, F., Cerling, C.E., Robinson, T.J., Douglas-Hamilton, I., 2006. Stable isotopes in elephant hair document migration patterns and diet changes. *Proc. Natl. Acad. Sci. USA* 103, 371–373.
- Chamaillé-Jammes, S., Fritz, H., Valeix, M., Murindagomo, F., Clobert, J., 2008. Resource variability, aggregation and direct density dependence in an open context: the local regulation of an African elephant population. *J. Anim. Ecol.* 77, 135–144.
- Chase, M.J., 2011. Dry Season Fixed-wing Aerial Survey of Elephants and Wildlife in Northern Botswana, September–November 2010. Tech. Rep. Elephants without Borders, Kasane.
- Codron, J., Codron, D., Lee-Thorp, J.A., Sponheimer, M., Kirkman, K., Duffy, K.J., Sealy, J., 2011. Landscape-scale feeding patterns of African elephant inferred from carbon isotope analysis of feces. *Oecologia* 165, 89–99.
- Delibes, M., Gaona, P., Ferreras, P., 2001. Effects of an attractive sink leading into maladaptive habitat selection. *Am. Nat.* 158, 277–285.
- Donovan, T.M., Thompson, F.R., 2001. Modeling the ecological trap hypothesis: a habitat and demographic analysis for migrant songbirds. *Ecol. Appl.* 11, 871–882.
- Douglas-Hamilton, I., Hillman, A.K.K., 1981. Elephant carcass and skeletons as indicators of population trends. In: Low-level Aerial Survey Techniques. Monograph 4. International Livestock Centre for Africa, Addis Ababa.
- Dudley, J., Criag, G., Gibson, D.S.C., Haynes, G., Klimowicz, J., 2001. Drought mortality of bush elephants in Hwange National Park, Zimbabwe. *Afr. J. Ecol.* 39, 187–194.
- Dunham, K.M., 2008. Detection of anthropogenic mortality in elephant *Loxodonta africana* populations: a long-term case study from the Sebungwe region of Zimbabwe. *Oryx* 42, 36–48.
- Dzialak, M.R., Webb, S.L., Harju, S.M., Winstead, J.B., Wondzell, J.J., Mudd, J.P., Hayden-Wing, L.D., 2011. The spatial pattern of demographic performance as a component of sustainable landscape management and planning. *Landscape Ecol.* 26, 775–790.
- Faluccci, A., Ciucci, P., Maiorano, L., Gentile, L., Boitani, L., 2009. Assessing habitat quality for conservation using an integrated occurrence-mortality model. *J. Appl. Ecol.* 46, 600–609.
- Gaughan, A.E., Stevens, F.R., Gibbes, C., Southworth, J., Binford, M.W., 2012. Linking vegetation response to seasonal precipitation in the Okavango–Kwando–Zambezi catchment of southern Africa. *Int. J. Remote Sens.* 33, 6783–6804.
- Guldemond, R.A.R., van Aarde, R.J., 2008. A meta-analysis of the impact of African elephants on savanna vegetation. *J. Wildlife Manage.* 72, 892–899.
- Hansen, M., DeFries, R., Townshend, J.R., Carroll, M., Dimiceli, C., Sohlberg, R., 2006. Vegetation Continuous Fields MOD44B, 2001 Percent Tree Cover, Collection 4. University of Maryland, College Park, Maryland.
- Harris, G.M., Russell, G.J., van Aarde, R.J., Pimm, S.L., 2008. Rules of habitat use by elephants *Loxodonta africana* in southern Africa: insights for regional management. *Oryx* 42, 66–75.
- Haynes, G., 1988. Longitudinal studies of African elephant death and bone deposits. *J. Archaeol. Sci.* 15, 131–157.
- Hebblewhite, M., Merrill, E.H., McDonald, T.L., 2005. Spatial decomposition of predation risk using resource selection functions: an example in a wolf–elk predator–prey system. *Oikos* 111, 101–111.
- Hoare, R.E., Du Toit, J.T., 1999. Coexistence between people and elephants in African savannas. *Conserv. Biol.* 13, 633–639.
- Illius, A.W., 2006. Linking functional responses and foraging behavior to population dynamics. In: Danell, K., Duncan, P., Bergstrom, R., Pastor, J. (Eds.), Large Herbivore Ecology, Ecosystem Dynamics, and Conservation. Cambridge University Press, Cambridge, pp. 71–96.
- Jarvis, A., Reuter, H.I., Nelson, A., Guevara, E., 2006. Hole-Filled Seamless SRTM Data V3. International Centre for Tropical Agriculture (CIAT). <<http://srtm.csi.cgiar.org>>.

- Johnson, C.J., Seip, D.R., Boyce, M.S., 2004. A quantitative approach to conservation planning: using resource selection functions to map the distribution of mountain caribou at multiple spatial scales. *J. Appl. Ecol.* 41, 238–251.
- Landscan, 2008. High Resolution Global Population Data Set. Copyrighted by UT-Battelle, LLC, operator of Oak Ridge National Laboratory under Contract No. DE-AC05-00OR22725 with the United States Department of Energy.
- Manly, B.F.J., Mcdonald, L.L., Thomas, D.L., Mcdonald, T.L., Erickson, W.P., 2002. *Resource Selection by Animals, Statistical Design and Analysis for Field*, second ed. Kluwer Academic Publishers, Dordrecht.
- McLoughlin, P.D., Gaillard, J.M., Boyce, M.S., Bonenfant, C., Messier, F., Duncan, P., Delorme, D., van Moorter, B., Saïd, S., Klein, F., 2007. Lifetime reproductive success and composition of the home range in a large herbivore. *Ecology* 88, 3192–3201.
- McLoughlin, P.D., Coulson, T., Clutton-brock, T., 2008. Cross-generational effects of habitat and density on life history in red deer. *Ecology* 89, 3317–3326.
- Metcalfe, S., Kepe, T., 2008. “Your elephant on our land”: the struggle to manage wildlife mobility on Zambian communal land in the Kavango–Zambezi transfrontier conservation area. *J. Environ. Dev.* 17, 99–117.
- Metzger, K.L., Sinclair, A.R.E., Hilborn, R., Hopcraft, J.G.C., Mduma, S.A.R., 2010. Evaluating the protection of wildlife in parks: the case of African buffalo in Serengeti. *Biodivers. Conserv.* 19, 3431–3444.
- Meyer, J.S., Irwin, L.L., Boyce, M.S., 1998. Influence of habitat abundance and fragmentation on northern spotted owls in western Oregon. *Wildlife Monogr.* 139, 3–51.
- Milner-Gulland, E.J., Bennett, E.L., 2003. Wild meat: the bigger picture. *Trends Ecol. Evol.* 18, 351–357.
- Moss, C.J., Croze, H., Lee, P.C., 2011. *The Amboseli Elephants: A Long-term Perspective on a Long-lived Mammal*. University of Chicago Press, Chicago.
- Nasseri, N.A., McBrayer, L.D., Schulte, B.A., 2011. The impact of tree modification by African elephant (*Loxodonta africana*) on herpetofaunal species richness in northern Tanzania. *Afr. J. Ecol.* 49, 133–140.
- Newmark, W.D., Manyaza, D.N., Gamassa, D.M., Sariko, H.I., 1994. The conflict between wildlife and local people living adjacent to protected areas in Tanzania: human density as a predictor. *Conserv. Biol.* 8, 249–255.
- Nielsen, S.E., Herrero, S., Boyce, M.S., Mace, R.D., Benn, B., Gibeau, M.L., Jevons, S., 2004. Modelling the spatial distribution of human-caused grizzly bear mortalities in the Central Rockies ecosystem of Canada. *Biol. Conserv.* 120, 101–113.
- Nielsen, S.E., Stenhouse, G.B., Boyce, M.S., 2006. A habitat-based framework for grizzly bear conservation in Alberta. *Biol. Conserv.* 130, 217–229.
- Nielsen, S.E., McDermid, G., Stenhouse, G.B., Boyce, M.S., 2010. Dynamic wildlife habitat models: seasonal foods and mortality risk predict occupancy-abundance and habitat selection in grizzly bears. *Biol. Conserv.* 143, 1623–1634.
- Norton-Griffiths, M., 1978. *Counting Animals*. African Wildlife Leadership Foundation, Nairobi.
- Owen-Smith, N., 2004. Functional heterogeneity in resources within landscapes and herbivore population dynamics. *Landscape Ecol.* 19, 761–771.
- R Core Team, 2012. *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0. <<http://www.R-project.org>>.
- Roever, C., van Aarde, R.J., Leggett, K., 2012. Functional responses in the habitat selection of a generalist mega-herbivore, the African savannah elephant. *Ecography* 35, 972–982.
- Roth, H.H., Douglas-Hamilton, I., 1991. Distribution and status of elephants in West Africa. *Mammalia* 55, 489–528.
- Tracks4Africa, 2010. Tracks4Africa Enterprises (Pty) Ltd. Unit 8 Innovation Center, Electron Street, Techno Park, Stellenbosch, 7599, Western Cape, South Africa. <<http://tracks4africa.co.za/>>.
- Trimble, M.J., Ferreira, S.M., van Aarde, R.J., 2009. Drivers of megaherbivore demographic fluctuations: inference from elephants. *J. Zool.* 279, 18–26.
- van Aarde, R.J., Jackson, T.P., 2007. Megaparks for metapopulations: addressing the causes of locally high elephant numbers in southern Africa. *Biol. Conserv.* 134, 289–297.
- van Aarde, R.J., Jackson, T.P., Ferreira, S.M., 2006. Conservation science and elephant management in southern Africa. *S. Afr. J. Sci.* 102, 1–4.
- van Aarde, R.J., Ferreira, S., Jackson, T., Page, B., De Beer, Y., Gough, K., Guldmond, R., Junker, J., Olivier, P., Ott, T., Trimble, M., 2008. Elephant population biology and ecology. In: Scholes, R.J., Mennell, K.G. (Eds.), *Elephant Management: A Scientific Assessment for South Africa*. Wits University Press, Johannesburg, pp. 84–145.
- Wall, J., Douglas-Hamilton, I., Vollrath, F., 2006. Elephants avoid costly mountaineering. *Curr. Biol.* 16, 527–529.
- Whyte, I., van Aarde, R.J., Pimm, S.L., 2003. Kruger’s elephant population: its size and consequences for ecosystem heterogeneity. In: Du Toit, J.T., Rodgers, W.A., Biggs, H.C. (Eds.), *The Kruger Experience: Ecology and Management of Savanna Heterogeneity*. Island Press, Washington, DC, pp. 332–348.
- Wittemyer, G., Elsen, P., Bean, W.T., Burton, A.C.O., Brashares, J.S., 2008. Accelerated human population growth at protected area edges. *Science* 321, 123–126.
- Wittemyer, G., Daballen, D., Douglas-Hamilton, I., 2013. Comparative demography of an at-risk african elephant population. *PLoS One* 8, e53726.
- Young, K.D., Ferreira, S.M., van Aarde, R.J., 2009. Elephant spatial use in wet and dry savannas of southern Africa. *J. Zool.* 278, 189–205.