

# Supporting conservation with biodiversity research in sub-Saharan Africa's human-modified landscapes

Morgan J. Trimble · Rudi J. van Aarde

Received: 24 January 2014 / Revised: 25 April 2014 / Accepted: 3 May 2014  
© Springer Science+Business Media Dordrecht 2014

**Abstract** Protected areas (PAs) cover 12 % of terrestrial sub-Saharan Africa. However, given the inherent inadequacies of these PAs to cater for all species in conjunction with the effects of climate change and human pressures on PAs, the future of biodiversity depends heavily on the 88 % of land that is unprotected. The study of biodiversity patterns and the processes that maintain them in human-modified landscapes can provide a valuable evidence base to support science-based policy-making that seeks to make land outside of PAs as amenable as possible for biodiversity persistence. We discuss the literature on biodiversity in sub-Saharan Africa's human-modified landscapes as it relates to four broad ecosystem categorizations (i.e. rangelands, tropical forest, the Cape Floristic Region, and the urban and rural built environment) within which we expect similar patterns of biodiversity persistence in relation to specific human land uses and land management actions. Available research demonstrates the potential contribution of biodiversity conservation in human-modified landscapes within all four ecosystem types and goes some way towards providing general conclusions that could support policy-making. Nonetheless, conservation success in human-modified landscapes is hampered by constraints requiring further scientific investment, e.g. deficiencies in the available research, uncertainties regarding implementation strategies, and difficulties of coexisting with biodiversity. However, information currently available can and should support efforts at individual, community, provincial, national, and international levels to support biodiversity conservation in human-modified landscapes.

---

Communicated by Karen E. Hodges.

---

**Electronic supplementary material** The online version of this article (doi:[10.1007/s10531-014-0716-4](https://doi.org/10.1007/s10531-014-0716-4)) contains supplementary material, which is available to authorized users.

---

M. J. Trimble (✉) · R. J. van Aarde  
Conservation Ecology Research Unit, Department of Zoology & Entomology, University of Pretoria,  
Private Bag X20, Hatfield, Pretoria 0028, South Africa  
e-mail: [morgantrimble@gmail.com](mailto:morgantrimble@gmail.com)

R. J. van Aarde  
e-mail: [rjvaarde@zoology.up.ac.za](mailto:rjvaarde@zoology.up.ac.za)

**Keywords** Cape Floristic Region · Countryside biogeography · Off-reserve conservation · Rangelands · Reconciliation ecology · Tropical forest

## Introduction

Conservation of biodiversity in Africa, like elsewhere, historically focused on the fortress model, whereby most protected areas (PAs) excluded people (see Carruthers 2009; Adams and Hulme 2001; Siurua 2006). Though PAs are essential for conservation success, they are insufficient (Rosenzweig 2003). For example, large mammal populations have been reduced by half in some African PAs since 1970 (Craigie et al. 2010), probably due, in part, to increasing isolation of PAs (Newmark 2008). Weak enforcement and ineffective management plague many of Africa's current PAs (Kiringe et al. 2007; Metzger et al. 2010; Pare et al. 2010), and many fail to cater to species with extensive spatial requirements, e.g. migratory animals (Holdo et al. 2010; Kirby et al. 2008; Thirgood et al. 2004; Western et al. 2009) and elephants *Loxodonta africana* (van Aarde and Jackson 2007). Even small-bodied species are not necessarily safe-guarded (Pauw 2007). Additionally, the configuration of PAs within the continent neglects key areas for biodiversity (Fjeldsa and Burgess 2008; Eardley et al. 2009; Chown et al. 2003; Fjeldsa et al. 2004; Beresford et al. 2011), a problem that may escalate if climate change makes PAs inhospitable to species they once protected (Loarie et al. 2009). If species' ranges shift with shifting climate, the areas crucial for their persistence will be transient (Hole et al. 2011). Furthermore, the scale of beta-diversity and habitat heterogeneity often extends far beyond that of individual PAs (Gardner et al. 2007), while human activities beyond PAs influence biodiversity within them (Hansen and DeFries 2007).

There are calls for an increased focus on biodiversity beyond African PAs (e.g. Eardley et al. 2009) on two fronts. First, conservation of some biodiversity elements depends on how well matrices outside of PAs cater for persistence. At the species level, for example, the Blue Crane *Anthropoides paradiseus* in South Africa (McCann et al. 2007), Ethiopia's critically endangered Sidamo lark *Heteromirafra sidamoensis* (Spottiswoode et al. 2009; Donald et al. 2010), and the last giraffes *Giraffa camelopardalis peralta* in West Africa (Ciofolo 1995) all depend on human-modified landscapes. At the ecosystem level, three biomes fall below the threshold 10 % protection status within the Afrotropic realm, i.e. tropical and subtropical dry broadleaf forests (6 %), montane grasslands and shrublands (8 %), and deserts and xeric shrublands (9 %), while several ecoregions are <5 % protected, especially when limited to IUCN categories I–IV, e.g. Southern Congolian forest-savanna mosaic (0 %) (Jenkins and Joppa 2009). Second, there are important links between biodiversity and ecosystem function, ecosystem services, and human livelihoods in working landscapes (Daily et al. 2001; Rosenzweig 2003). For example, maintaining natural habitat in and around farms can enhance pollination and, thus, has an economic value to production landscapes (Carvalho et al. 2010; Munyuli 2012), and natural systems in Africa provide economic and nutritional benefits to both rural and urban dwellers (Tabuti et al. 2009; Schreckenber 1999; Vanderpost 2006).

Though, scientists have neglected the biogeography of human-modified landscapes in sub-Saharan Africa, ecologists are increasingly studying the capacity of such landscapes to support biodiversity (Trimble and van Aarde 2012). Such studies are required in order for policy-makers to make defensible decisions regarding land use in relation to biodiversity

conservation goals in the face of rapid economic development that could potentially decimate biodiversity. Agriculture in Africa has been characterized by traditional, labor-intensive, smallholder enterprise; production has been low and has remained relatively stagnant (Abate et al. 2000; Deininger et al. 2011). However, economic development and population growth are driving change in African landscapes, with several nations among the world's fastest growing economies (IMF 2013). In 2009, the population reached one billion and is predicted to double by 2050 (UN-HABITAT 2010). Urbanization is a strong force; 40 % of the current African population is city-dwelling, and by 2050, 60 % will be urban (UN-HABITAT 2010). Even so, the rural population will also grow substantially, predicted to increase by nearly 50 % by mid-century (UN Population Division 2012), while growing urban centers will depend heavily on rural resources. To meet this demand and to improve food security, there are calls for both intensifying smallholder agriculture (Baiphethi and Jacobs 2009; Snapp et al. 2010; Muriuki et al. 2005; Baudron et al. 2011) and extensifying production (Muriuki et al. 2005).

Therefore, the interest in biodiversity in human-modified lands is timely. Although Africa's natural ecosystems are more intact than many other regions', a proactive approach to biodiversity conservation that strives for the most prudent management of the unprotected matrices between PAs is clearly preferable to trying to reconnect and restore already degraded ecosystems (Gardner et al. 2010). Thus, as policy-makers chart the future course of development in Africa, they should consider the effects of different choices on biodiversity in human-modified lands, what steps can be taken to prevent biodiversity loss, and the benefits and costs of biodiversity persistence to people. Failure to do so may necessitate expensive restoration and reintroduction efforts. Studies of biodiversity patterns and the processes that maintain them in human-modified landscapes provide an evidence base to support defensible management that meets the needs of people and biodiversity simultaneously. The evidence base should, furthermore, provide for relevant ecological contexts. For example, management standards for timber plantations aim to minimize impact on biodiversity in surrounding natural forests. Yet, the same standards have been applied in plantations embedded in grasslands with dubious efficacy for minimizing impacts on grassland biodiversity (Lipsey and Hockey 2010; Pryke and Samways 2003).

This scientific focus on biodiversity in human-modified landscapes is distinct from Africa's thirty-some-year experiment in community-based conservation (CBC, but also known as Integrated Conservation and Development Projects, Community-Based Natural Resource Management, and others), but these two fields can and should be amalgamated. Promoters of CBC claim that it increases the chance of conservation success and simultaneously reduces rural poverty by allowing community involvement in management and profit from natural resources, especially large mammals (see Hackel 1999). The philosophy of linking wildlife conservation and rural economic development and the practical successes and failures therein have been discussed in a large body of literature (e.g. Songorwa et al. 2000; Torquebiau and Taylor 2009; Hackel 1999) and are subjects of renewed controversy outside of Africa (see Kareiva and Marvier 2012; Doak et al. 2013). However, the discussion has focused on socioeconomics and politics with fleeting consideration for assessing actual biodiversity persistence under different CBC models, a problem pointed out by Caro (1999) and subsequently largely ignored.

In this review, our objective was to elucidate the state of knowledge regarding biodiversity in sub-Saharan Africa's human-modified landscapes to provide an overview for scientists and policy-makers. We also sought to identify constraints and opportunities for integrating biodiversity conservation with land-use management that require further attention in order to support the implementation of sensible policies.

## Methods

### Literature search

We searched the ISI Web of Knowledge (up to 2012) with keywords “Africa” and “biodiversity or conservation” and each of the following terms: “agricultur\*”, “agroforestry”, “communal”, “farm\*”, “game farm”, “game ranch”, “human-modified”, “multiple-use management”, “periurban”, “private nature reserve”, “rangeland”, “rural”, “suburban”, and “urban”. We also searched for the terms “countryside biography”, “reconciliation ecology”, and “off-reserve conservation”. Additionally, we included relevant papers found coincidentally or in reference lists.


### Biodiversity in human-modified landscapes of African ecosystems

In summarizing the literature on biodiversity in Africa’s human-modified landscapes, we separate our discussion into four major ecosystem types (see Fig. 1) within which we expect similar patterns to emerge. (1) Rangelands attract the bulk of our attention as Africa’s biggest ecosystem type, and rangeland biodiversity is perhaps the most compatible with human land-uses, so biodiversity-conscious land-use planning in rangelands could yield huge benefits. (2) Tropical forests are discussed briefly with a focus on Central and East African forests, and we refer readers to an excellent review of the abundant literature from West Africa (Norris et al. 2010). (3) The Cape Floristic Region, though small, is extremely rich in species yet threatened by extensive commercial development, and we discuss a growing body of literature on land-use management in the region. Finally, (4) the urban and rural built environment will become an increasingly important concern for biodiversity conservation in Africa where the increase of urban land cover is predicted to be the highest in the world at nearly 600 % in the first three decades of the twenty-first century (Seto et al. 2012); proper management and infrastructure development could attenuate the consequences for biodiversity.

### Rangelands

Two-thirds of sub-Saharan Africa is composed of rangelands (Fig. 1), consisting of arid and semi-arid grasslands, woodlands, savannas, shrublands, and deserts. The rural people inhabiting rangelands are typically agropastoralists, specializing in small-scale farming or livestock keeping or a combination. Some agricultural practices in rangelands may be harmful to biodiversity, e.g. overcultivation, overgrazing (Kerley et al. 1995), bush fires, cultivation of marginal and easily eroded land, and widespread use of chemicals and pesticides (Darkoh 2003). Many people in rangelands depend heavily on wild resources, e.g. via hunting and gathering or by profiting from wildlife tourism (Homewood 2004). Game ranching is an increasingly popular land-use option across African rangelands (McGranahan 2008), and so are “eco-estates” (Grey-Ross et al. 2009b), where people choose to live amongst the natural beauty of African rangelands and their considerable species diversity, especially charismatic large mammals.

The ecological mechanisms that maintain different rangeland types in different locations, e.g. grassland versus woodland, are not fully understood though interactions between soils, climate, fire, herbivory, and human disturbance are thought to be important (see Bond and Parr 2010). The biggest threats to grasslands include afforestation or bush



**Fig. 1** Map of sub-Saharan Africa showing ecosystem types adapted from Olson et al. (2001): rangelands (desert and xeric shrubland, montane grassland/shrubland, flooded grassland/savanna, and tropical/subtropical grassland/savanna/shrubland), tropical forests (moist and dry tropical forest), the Cape Floristic Region (Mediterranean forest/woodland/scrub), and the urban and rural built environment represented by the human influence index (Wildlife Conservation Society and Center for International Earth Science Information Network 2005), a dataset comprising nine data layers incorporating population pressure (population density), human land use and infrastructure (built-up areas, nighttime lights, land use, land cover), and human access (coastlines, roads, railroads, navigable rivers)

encroachment and clearing for agriculture (Bond and Parr 2010), while threats to the woodlands include woodcutting, clearing for agriculture, and over-use (Tabuti 2007; Schreckenber 1999). Many perceive that biodiversity is declining in rangeland systems; they blame poor agricultural practices, land conversion, and over-utilization of wild resources by rural people and worry that these patterns will increase with population growth (e.g. Darkoh 2003; Thiollay 2006). However, documented evidence of biodiversity loss in rural rangelands is sparse. Many areas have likely lost some species, but surprisingly, long-inhabited regions lacking formal PAs, e.g. Kenya's Laikipia district, maintain abundant wildlife including large carnivores and elephants (Gadd 2005; Kinnaird and O'Brien 2012) that might seem at odds with human occupation (Woodroffe et al. 2007). Rangeland systems are characterized by disturbances such as fire, unpredictable rainfall, grazing and browsing pressure, and physical disturbance. Therefore, rangeland biodiversity may be relatively resilient to anthropogenic disturbance due to the ability to disperse, colonize, and persist in patchy, fluctuating environments (Homewood 2004). Thus, human-modified landscapes have the potential to maintain a relatively large portion of rangeland biodiversity.

Nonetheless, conservation in rangelands has traditionally excluded people from designated PAs. In South Africa, for example, conservation planning often dichotomizes ‘human land-use’ and ‘conservation’ with little consideration for different land-use options that may be variably amenable to biodiversity (e.g. Wessels et al. 2003; Chown et al. 2003). On the other hand, some authors have called to “mainstream” conservation into human-modified lands (e.g. Soderstrom et al. 2003; Pote et al. 2006). O’Connor and Kuyler (2009) used expert opinion to rank the impact of land uses in moist grasslands on overall biodiversity integrity (from least to most impact: conservation, game farming, livestock, tourism, crops, rural, dairy, timber, and urban). Empirical studies are amassing to assess such assertions, which could support land-use planning for conservation. Here we discuss emerging research on biodiversity in several of the most common rangeland land uses.

### *Grazing*

Grazing is important to the maintenance of grassland and savanna habitats, economic development, and management for biodiversity. However, plant responses to grazing are idiosyncratic and incompletely understood (see Watkinson and Ormerod 2001; Rutherford et al. 2012). Overgrazing can lead to degradation and bush encroachment (the proliferation of woody plants at the expense of grasses), while too little grazing can result in succession to woodland (Watkinson and Ormerod 2001). Of course, grazing effects on vegetation can affect higher trophic levels as well, so it is important to understand vegetation responses to grazing, not only for livestock production, but also because vegetation dynamics affect many other species. However, not all grazing landscapes are alike; unique vegetation dynamics in different ecosystems mean that landscapes can respond disparately to grazing pressure (Todd and Hoffman 2009).

Research is emerging that investigates aspects of grazing management and biodiversity in Africa; we summarize 30 such studies in Online Resource Table 1. Generally, these studies look at grazing intensity, or proxies such as bush encroachment, and show that many wild species may be maintained depending on management and location. For example, traditional pastoral practices, i.e. burning and boma creation, may even be necessary to maintain avian diversity in some East African savanna areas (Gregory et al. 2010). Contrastingly, bush encroachment due to overgrazing in Ethiopia may provoke Africa’s first avian extinction (Donald et al. 2010; Spottiswoode et al. 2009).

Online Resource Table 1 shows that only about a third of studies compared biodiversity of livestock grazing landscapes to controls with indigenous grazers such as PAs. Most studies came from South Africa (67 %), and most assessed grazing effects on plants (43 %) or insects (27 %). Many areas of investigation remain open, such as the role of vegetation structure including keystone, isolated trees in maintaining biodiversity in human land-use areas; such trees are important for maintaining diversity in natural systems (Dean et al. 1999). A common conclusion with regards to plant diversity is that spatial heterogeneity in grazing management that includes PAs will enhance gamma diversity because different species thrive at different grazing intensities (e.g. Fabricius et al. 2003).

### *Agricultural mosaic*

While extensive grazing is common in arid-savannas and xeric shrublands, a land-use mosaic of grazing and cropping interspersed with settlements is common in more mesic savannas and grasslands. This mosaic effect may have important consequences for the maintenance of biodiversity, and studies of biodiversity in agricultural mosaics (24 studies

summarized in Online Resource Table 2) identify some common themes. Compared to strict PAs, agricultural mosaics may actually be beneficial to some species groups. For example, Caro (2001) illustrated greater diversity and abundance of the small mammal assemblage in the agricultural matrix outside Katavi National Park, Tanzania than inside, a pattern also found for Niokolo Koba National Park, Senegal (Konecny et al. 2010). Richness of birds, amphibians, small mammals, butterflies, and trees is similar at 41 sites across a land-use gradient from Katavi National Park to non-intensive agricultural land; however, composition changes along the gradient, and although the PA holds some unique species, some species found outside the PA are absent within (Gardner et al. 2007). Thus, agricultural mosaics may contribute to greater gamma diversity at the landscape scale; nonetheless understanding the conservation implications of higher gamma diversity may require a regional or global perspective on species rarity and commonness.

It is a common finding that agricultural intensification (e.g. mechanization, shortening fallows, destruction of remnant habitat patches, and introduction of cash crops) can have detrimental effects on the biodiversity value of agricultural mosaics. The mosaic effect of traditionally managed farms in KwaZulu-Natal, South Africa may support, and even enhance, bird diversity (Ratcliffe and Crowe 2001), but intensification results in species declines due to loss of 'edge' habitats. In Burkina Faso, common butterfly species occur in cultivated areas, while specialists are more common in old fallows and grazed areas, probably because grazing maintains host plants and, thus, diversity (Gardiner et al. 2005). In this case, an agricultural mosaic of shifting fallows could support butterfly meta-populations that allow species persistence, while intensification could be detrimental (Gardiner et al. 2005). In Ethiopian grasslands, low-intensity agriculture supports moderate plant diversity, while larger-scale, mechanized farms reduce tree cover and diversity (Reid et al. 1997). Similarly, in the Serengeti-Mara ecosystem, commercial mechanized agriculture is associated with declining wildlife populations (Homewood et al. 2001; Homewood 2004).

### *Cropping*

Cropping is perhaps more at odds with biodiversity than grazing is because cropping involves the direct removal of indigenous vegetation and planting of, generally, non-indigenous species. Nonetheless, crops can support wild species, and conservation value may depend on the crops planted, the farming methods employed, and the arrangement of fields with respect to natural habitat. Relatively few studies assessed biodiversity solely in cultivated areas (10 studies summarized in Online Resource Table 3), as opposed to agricultural mosaics (Online Resource Table 2). This perhaps reflects the current state of African agriculture, where most farms are smallholder or subsistence based rather than expansive, commercial cultivation. Although there are exceptions, average farm size is just 2–3 ha (Deininger et al. 2011). Where commercial cultivation does occur, loss of biodiversity may be seen as a foregone conclusion not worth investigating (see Thiollay 2006). Many studies of biodiversity in cultivation were concerned primarily with the benefits of that diversity for production via pest control, fertility enhancement, or pollination services, rather than for its value to conservation (e.g. Carvalheiro et al. 2010; Midega et al. 2008; Tchabi et al. 2008).

### *Agroforestry*

Agroforestry, the integration of trees into agriculturally productive landscapes, has garnered much attention in the global conservation community because it has been shown to

provide habitat for relatively high levels of forest species diversity (see Bhagwat et al. 2008). In African rangelands, agroforestry can be divided into two types: technological and traditional. Technological agroforestry deals with the expertise to plant and maintain tree species that increase productivity in agricultural production systems. Kenyan farmers, for example, plant crops of fodder trees, which raise milk yields of cows and goats (Pye-Smith 2010a). Government programs in Niger, Zambia, Malawi, and Burkina Faso support large-scale ‘evergreen agriculture’ projects to plant indigenous trees such as *Faidherbia albida* among crops, which maintain green cover year-round, increase yields by improving soil fertility, and provide fodder and firewood (Garrity et al. 2010). Evergreen agriculture and other technological agroforestry projects are touted as having greater biodiversity value than do monoculture crops (see Pye-Smith 2010a, b; Kalaba et al. 2010; Garrity et al. 2010). Yet, evidence to support these claims remains mostly anecdotal, warranting further research because plans are underway to expand technological agroforestry projects throughout Africa (Garrity et al. 2010).

Traditional agroforestry, on the other hand, is a millennia-old practice, particularly evident in the parkland savannas of West Africa, of people maintaining savanna tree species in pastures, fields, and villages. These trees provide shade, food, wood, and even cash when commercially traded (e.g. shea, baobab), and traditional agroforestry may contribute to the maintenance of tree species in addition to species for which trees provide habitat. Many studies have enumerated tree diversity in farmlands (Online Resource Table 4). Even so, the conservation value of agroforestry varies. Augusseau et al. (2006) report that in Burkina Faso, few indigenous species are important to farmers and none are planted. Even where tree richness is maintained at a relatively high level, the persistence of trees in traditional agroforestry can be compromised if the economic value of totally clearing the land, e.g. for mechanized agriculture or firewood, outpaces the value of non-timber products (Tabuti et al. 2009). Additionally, based on demographic profiles of tree species, tree regeneration appears problematic in many human-modified landscapes (e.g. Fandohan et al. 2010; Schumann et al. 2010; Venter and Witkowski 2010). For example, a study in Benin shows that the largest shea trees are often in villages or fields, but seedling survival is low compared to nearby PAs (Djossa et al. 2008). Regeneration potential can also be diminished when harvesting tree products affects recruitment, as is the case for *Khaya senegalensis* in Benin (Gaoue and Ticktin 2008). Where natural regeneration potential is compromised, intervention may be required to ensure rejuvenation (Kindt et al. 2008; Ouinsavi and Sokpon 2008), especially if farmers abandon traditional rotational land-use systems such as long fallow, where trees are often most capable of regenerating (Raebild et al. 2007; Schreckenberg 1999).

Fortunately, agroforestry management in rangeland ecosystems is an active area of research with regards to developing strategies to encourage tree persistence (Kindt et al. 2008; Tabuti et al. 2009; Augusseau et al. 2006). Yet, there is a surprising lack of research to assess the value of savanna agroforests for faunal diversity or even non-tree plant diversity (Online Resource Table 4), aspects that have been more thoroughly studied in the tropical forest context (Bhagwat et al. 2008), and this dearth should be remedied.

### *Game ranching and private nature reserves*

The wildlife industry, including game ranching, game farming, and private nature reserves, has become big business, especially in southern and East African rangelands. These land-use options involve profiting from consumptive (e.g. trophy hunting, live animal sales, meat) or non-consumptive (e.g. tourism, aesthetic value) use of wildlife on communal or

private land. South Africa alone has an estimated 9,000 private game ranches, covering 20.5 million ha, many of which were converted from traditional livestock ranches (NAMC 2006). Ranching game rather than domestic livestock may ameliorate effects of overgrazing because indigenous species have coevolved with indigenous vegetation (Kerley et al. 1995), and indigenous browsers may control bush encroachment (McGranahan 2008; Taylor and Walker 1978). Thus, the wildlife industry may be a boon to biodiversity conservation; however, very few studies have actually assessed impacts on biodiversity, which may be positive or negative and likely depend on management actions (Cousins et al. 2008).

Occurrence and abundance of mammal species on private land has increased due to game ranching (Lindsey et al. 2009). Nonetheless, some aspects of the wildlife industry are worrying. Privatization of wildlife (and sometimes legislative requirements) begets ubiquitous game fencing (McGranahan 2008; Lindsey et al. 2009) with substantial ecological consequences including the interruption of natural movements, inbreeding, and overstocking (Lindsey et al. 2009; Hayward and Kerley 2009). Ranches are often quite small (South African provincial averages range from 8.2 to 49.2 km<sup>2</sup>), and smaller ranches necessitate more intensive management interventions (Lindsey et al. 2009; Bothma 2002). Additionally, the industry's focus on trophies may skew natural communities in favor of valuable species and induce semi-domestication (Mysterud 2010), and it has resulted in extra-limital introductions, questionable breeding practices, and persecution of predators (Lindsey et al. 2009). Even within the mammal community, generally the focus of game ranching, the full complement of species of a given ecosystem may not be maintained on ranches despite deliberate re-introductions (Grey-Ross et al. 2009a).

Thus, much more research is needed on the biodiversity value of the wildlife industry and what measures, e.g. promoting conservancies over single game ranches (Lindsey et al. 2009), can improve this value. Best-practice management in terms of grazing pressure, fire regimes, bush encroachment, wildlife ownership policies, and fencing needs more attention (McGranahan 2008). Furthermore, surprisingly little is known about the impacts of game ranching on species other than large mammals. Even so, game ranches are likely more amenable to most indigenous biodiversity than are many other commercial land-use options. For example, large eagles in South Africa's Karoo shrublands are much more common in areas stocking indigenous mammals than in areas with domestic livestock and cultivation (Machange et al. 2005).

### Tropical forests

Though rangelands cover the majority of Africa, tropical forests also make up a considerable portion [ $\sim 20\%$  (Brink and Eva 2009), Fig. 1], particularly rich in biodiversity. Research on biodiversity in human-modified landscapes is biased towards tropical forests (Trimble and van Aarde 2012). Nonetheless, biodiversity in human-modified tropical forest landscapes in Africa has received much less scientific attention than in other regions, especially South and Central America (Gardner et al. 2010). African tropical forests tend to be in less conflict with high human population densities than elsewhere (e.g. Southeast Asia and Brazilian Atlantic forests) (Gardner et al. 2010), although in West Africa 80% of the original forest extent is now an agricultural-forest mosaic home to 200 million people (Norris et al. 2010).

We do not attempt a comprehensive review of African tropical forest biodiversity in human-modified landscapes and refer readers to Norris et al. (2010) for an excellent treatment of the West African scenario. They lament the lack of data regarding biodiversity

in African agricultural-forest mosaics but are able to reach some general conclusions. Land uses that maintain tree cover are more amenable to forest biodiversity than those that do not. Species richness increases in some modified habitats, such as logged and secondary forest, for some species groups, but endemic forest species are often lost. Additionally, relatively high species richness in modified habitats comprises, in part, species not present in the baseline forest comparison, so species richness alone likely overestimates the value of modified habitats for forest species. Furthermore, habitat modification seems to affect richness of forest plant species more negatively than of some animal groups.

Although logically, it seems more difficult to encourage the persistence of biodiversity in human-modified landscapes embedded in tropical forests than in rangelands, research can indicate best practices for land-use planning. In contrast to West Africa, Central Africa still maintains large tracts of relatively undisturbed forest that are becoming increasingly threatened by development, and lessons from studying African forest biodiversity in human-modified landscapes should be incorporated into development policy for the region (Norris et al. 2010).

The tropical forest biome extends to East and southern Africa where forests are less extensive, confined largely to high altitudes inland and a linear belt along the coast. These geographic constraints present unique challenges for conservation and heighten the importance of maintaining endemic species and retaining connectivity in fragmented forests. Fewer studies consider East and southern African tropical forests than West African forests, but work is emerging to support land-use planning in the region, and results largely conform to those found for West Africa. Agroforestry in Ethiopian and Tanzania supports less diversity than forests but more than other land uses (Gove et al. 2008; Hemp 2006; Hall et al. 2011; Negash et al. 2012). While Schmitt et al. (2010) found higher overall plant richness in Ethiopian coffee agroforests than natural forests, richness of typical forest species was lower. In Kenya, connectivity of coastal forest fragments for primates may be influenced by matrix structure (Anderson et al. 2007). Farmland outside tropical forest remnants, especially structurally complex subsistence farms, support higher bird richness than forests; however, many forest species are lost, highlighting the importance of maintaining the forest remnants but also supporting traditional farming techniques over commercial monocultures (Laube et al. 2008; Mulwa et al. 2012). Furthermore, structurally diverse farmland surrounding forest remnants may enhance forest pollinator communities (Hagen and Kraemer 2010). Similarly, South African forest remnants embedded in various matrix types have similar bird species richness, but abundance is highest in fragments in agricultural matrices due to the presence of forest generalists and open-habitat species, while forest specialists are rare (Neuschulz et al. 2011). Additionally, herpetofaunal richness does not decline monotonically along a land-use gradient from forest to cultivation, while richness of functional groups erodes along the gradient due to sensitivity of some specialist groups (Trimble and van Aarde 2014). Forest fragments and grasslands in the agricultural mosaic outside a PA in southern Mozambique have more beetle species and higher abundance, while endemic beetle species are better represented inside the PA (Jacobs et al. 2010).

### Cape Floristic Region

While small in area (approximately 90,000 km<sup>2</sup>, see Fig. 1), the Cape Floristic Region (CFR) of South Africa is a biodiversity hotspot of global significance (Myers et al. 2000) consisting of a Mediterranean-type ecosystem with high species turnover across the landscape and high endemism. Systematic conservation planning has been conducted for

the region but focused on pristine habitat that could be formally protected (see Cowling and Pressey 2003). Because spatial turnover of species is so high, however, successful conservation will depend heavily on efforts in human-modified landscapes beyond PAs (Cox and Underwood 2011). Based on species-area curves for plants and vertebrates in the CFR, practicing biodiversity friendly management on just 25 % of the land that is beyond PAs, but still in a natural or semi-natural state, might add an additional 541 species to the 7,340 estimated to occur in PAs (Cox and Underwood 2011).

However, in contrast to many areas of Africa dominated by subsistence agriculture, the CFR is characterized by large areas of intensively managed agricultural monocultures with low biodiversity value (Giliomee 2006). Overall, only 26 % of the CFR has been transformed, but the CFR is made up of different habitat types, and some, especially in the fertile lowlands, have lost much more of their area to cultivation, urbanization, and heavy invasion of exotic plants; for example, coast renosterveld is more than 80 % transformed (Rouget et al. 2003a). Transformation threatens not only the CFR's plants but also endemic and vulnerable animals such as the Black Harrier *Circus Maurus*, which has been displaced from the inland plains by cereal agriculture and now breeds, less successfully, in the coastal strip and inland mountain habitats (Curtis et al. 2004). Though the Black Harrier can forage in cultivated areas, it relies on intact vegetation to breed (Curtis et al. 2004).

PAs within the CFR are concentrated in areas of low agricultural value (e.g. mountains and coastlines), so biodiversity in fertile areas depends on conservation on privately owned land (Giliomee 2006; Rouget et al. 2003b). To increase the biodiversity value of agricultural areas, the primary focus should be on conserving remnants of natural vegetation on farms (Giliomee 2006). This is being attempted through incentive-driven stewardship agreements that protected almost 70,000 ha of vegetation on private land between 2003 and 2007 (Von Hase et al. 2010). Additionally, farm management practices may be variably amenable to biodiversity. For example, though vineyards have very different arthropod communities than those in natural vegetation, organic vineyards support greater diversity than do more intensively managed vineyards (Gaigher and Samways 2010). However, these effects may be taxon dependent; for instance, organic vineyard management benefits richness of monkey beetles (crucial pollinators), but not bees (Kehinde and Samways 2012). Similarly, apple orchards support less arthropod diversity than natural vegetation does, but orchards that are not sprayed with pesticides have a higher diversity than sprayed sites (Witt and Samways 2004). On the other hand, farms with a mixture of different crops and remnants of natural vegetation maintain most fynbos bird species and attract several additional species, while single crop sites without remnant vegetation have much less bird diversity and lose many fynbos species (Mangnall and Crowe 2003). Clearly, maintaining remnant vegetation and connectivity in agricultural areas of the CFR is crucial, but more research is needed to tailor agricultural practices to better conserve CFR species in production landscapes.

### Urban and rural built environment

Plant and vertebrate species richness and endemism are correlated with human population density and human infrastructure in sub-Saharan Africa (Burgess et al. 2007; Fjeldsa and Burgess 2008; Balmford et al. 2001), which is substantial in many regions (see Fig. 1). That the pattern endures in relatively developed South Africa means either that species persist to some degree with humans in disturbed habitats at current levels, that human-disturbed habitats actually attract more species, or that a major extinction debt is yet to be paid (Fairbanks 2004; Chown et al. 2003). Regardless, areas with high human density

[predicted to increase dramatically in Africa, outpacing growth in all other regions in the coming decades (Seto et al. 2012)] require appropriate regulations to ensure they remain as amenable as possible to biodiversity conservation. This will be especially important in Africa's most biologically rich yet rapidly urbanizing regions; by 2030 for example, the urban area within the Eastern Afromontane and Guinean Forests of West Africa hotspots is forecasted to be 1,900 and 920 % of 2,000 levels respectively (Seto et al. 2012).

Some obvious steps include discouraging urban sprawl; providing appropriate housing for low income populations while controlling illegal settlements in biodiversity sensitive areas; designing relevant green spaces that include aquatic habitats and indigenous plants; and managing invasive species, waste, and pollutants (Puppim de Oliveira et al. 2011; Muriuki et al. 2011). Research on managing Africa's urban and rural built environments for biodiversity is in its infancy and is mostly constrained to South Africa. Clearly, more research is needed, yet several studies provide pertinent information for planners.

While urban environments might not seem particularly hospitable to biodiversity, even small home gardens in African cities can harbor a remarkable number of species, especially in the tropics, both intentionally cultivated and otherwise (Cumming and Wesolowska 2004; Lubbe et al. 2010; Bigirimana et al. 2012). In South Africa, socioeconomics, urbanicity, and ecological factors influence plant diversity and the proportion of invasive species in home gardens (Lubbe et al. 2010; Molebatsi et al. 2010). Gardens with a high number of non-indigenous species contribute to biotic homogenization and pose the risk of new introductions that could prove detrimental to indigenous ecosystems. Therefore, invasive species in the urban landscape need to be controlled through regulation and removal, especially in threatened and fragile ecosystems (Alston and Richardson 2006; Dures and Cumming 2010; Cilliers et al. 2008; Bigirimana et al. 2012).

Green spaces such as city parks, tree-lined streets, and even golf courses in urban environments can support certain species. Dures and Cumming (2010) show that habitat quality, rather than patch metrics such as area, have the greatest influence on bird diversity in fynbos in an urban gradient in Cape Town. Thus, controlling invasive species even in high-density housing areas may be more beneficial for birds than expanding the low quality network of urban reserves. Alien pine tree removal helps restore invertebrate species diversity in Cape Town, and fragments of natural vegetation and gardens with indigenous plants help maintain it (Pryke and Samways 2009). In the Durban Metropolitan Open Space System, complex habitats (i.e. with trees and shrubs) support higher invertebrate diversity than simplified habitats (i.e. mown lawns); however, simple habitats might cater for certain rare species (Whitmore et al. 2002). Green spaces in urban Pretoria contribute to butterfly and moth diversity (McGeoch and Chown 1997) and also support indigenous birds (van Rensburg et al. 2009), while maintaining urban riparian vegetation is necessary for dragonfly conservation in Pietermaritzburg (Samways and Steytler 1996). Better ecological planning for developments such as golf courses could increase the likelihood for biodiversity persistence and minimize negative consequences, even in the CFR (Fox and Hockey 2007). Additionally, habitat engineering, e.g. creating biotopes for dragonflies (Steytler and Samways 1995), might be a useful tool in the urban context to promote biodiversity, although continual management of these habitats may be necessary to ensure species persistence (Suh and Samways 2005).

When species are range-restricted such that a single metropolitan area may affect most of their range, special attention is required. For example, two small forest parks in Durban suburbs are home to the last remnant populations of the rare tree *Oxyanthus pyriformis* whose specialist pollinators, the long-tongued hawkmoths, appear unable to tolerate sub-urban living. Hand pollination and planting of seedlings will be necessary to maintain the

species (Johnson et al. 2004). Similarly, conservation of plants in Cape Town is hampered by apparent sensitivity of specialist pollinator birds to urbanization, which is concerning given the increasing urbanization in the CFR (Seto et al. 2012). Durban covers a large portion of the range of the black-headed dwarf chameleon *Bradypodion melanocephalum*, and translocations from sites demarcated for development to sites reserved for conservation have proven somewhat successful, dependent on adequate alien-plant-control and habitat restoration (Armstrong 2008). Unique landscape features within urban areas may also require special attention. For example, Table Mountain in Cape Town harbors endemic species whose conservation depends not only on the PA of Table Mountain but also on management of lower elevation suburban woodlands (Pryke and Samways 2010).

On the rural end of the settlement spectrum, less attention has been given to biodiversity persistence. Some agricultural mosaic studies consider rural settlements, but a few studies treat it explicitly. For example, similar to shifting cultivation, some cultures practice shifting settlement, and abandoned settlements have been shown to provide valuable seasonal resources, e.g. fruit trees, to chimpanzees *Pan troglodytes* in Mali (Duvall 2008). Even road verges may provide for some types of biodiversity. For example, verges in the Karoo support some plant species not found in adjacent grazing lands, though many species from pastures are not found in verges (O'Farrell and Milton 2006). Verges also support invertebrates and could prove valuable to conservation because verges are public spaces that can be managed for biodiversity (Tshiguvho et al. 1999).

Understanding more about urban settlement and biodiversity may even benefit conservation in once remote PAs where rural sprawl and infrastructure for wildlife tourism can be dramatic (Wittemyer et al. 2008). For example, recent decades have seen substantial increases in rural sprawl along with the construction of 60 tourist lodges, 1,200 boreholes, and 540 km of roads in the Okavango Delta, one of Botswana's premiere conservation areas (Vanderpost 2006).

## Constraints and opportunities

The scope of this review was broad, covering four major ecosystem types within sub-Saharan Africa. Specific conclusions are difficult because species' responses to land use are clearly idiosyncratic and dependent on local factors. Nonetheless, we have identified some general conclusions and constraints summarized in Table 1 and discussed below.

### The science of biodiversity in human-modified landscapes

As others have pointed out, understanding the value of human-modified landscapes for biodiversity, especially in Africa, is hampered by data constraints (Norris et al. 2010; Pettoirelli et al. 2010; Waltert et al. 2011; Trimble and van Aarde 2012). Many studies are limited in temporal and spatial scale, and poor study design may result in insufficient sampling of habitats. The focus on species richness of certain habitat types while failing to account for the importance of species from other habitats in assigning conservation value to different land-use options may neglect the bigger picture; Bond and Parr (2010), for example, call for more collaboration between forest conservationists and others. More consideration for the value of different species in terms of commonness and rarity also needs to be developed because human-modified landscapes often fail to cater for endemic and specialist species (Waltert et al. 2011), and a better understanding of beta and gamma diversity at a landscape scale is necessary.

**Table 1** General conclusions regarding practices that support biodiversity in human-modified landscapes and scientific and implementation concerns requiring further investigation

---

**Practices that tend to support species diversity and richness in human-modified landscapes**

- Prefer diversity in selection of crops grown (i.e. polyculture) and land use (i.e. land-use mosaics, over homogenous monocultures)
- Encourage traditional agricultural practices over large-scale, mechanized farming
- Leave as much remnant natural vegetation as possible and monitor or assist maintenance of keystone structures or species, e.g. large trees
- Ensure strict protection for specialist and endemic species and expand PA coverage focused on these groups
- Encourage appreciation and understanding of conservation goals among land users
- Discourage urban sprawl and maintain and manage urban green spaces
- Favor use of native species in gardens and cultivation

**Avenues for further investigation into scientific uncertainties and implementation practices**

- Researching poorly documented combinations of species group, ecosystem type, and land use, e.g. mammals in rangeland agroforests
  - Moving beyond occurrence data to likelihood of persistence, e.g. how dependent are species in human-modified landscapes on nearby PAs or remnant habitat?
  - Investigating the value of reintroduction or rewilding in human-modified landscapes
  - Going beyond the species level of biodiversity to integrate genetic and ecosystem concepts
  - Supporting technological and traditional knowledge for cultivating useful native species and investigating the effects of these practices on other taxa
  - Creating frameworks for valuing importance of different species in different landscapes at a local level within a global context, e.g. commonness versus rarity and specialist versus generalist
  - Developing policies that integrate and account for local, regional, and global conservation needs and land use systems
- 

Additionally, further investigation into the relationship between occurrence and persistence is required, as are more studies that delve beyond species richness into the processes that support the observed patterns of biodiversity. For example, studies of demographic processes (e.g. Venter and Witkowski 2010; Djossa et al. 2008; Schumann et al. 2010) and population trends (e.g. Trimble and van Aarde 2011; Stoner et al. 2007) for species inhabiting human-modified landscapes can provide insight beyond mere patterns of occurrence. Furthermore, as elsewhere (Gardner et al. 2010), studies of biodiversity in African human-modified landscapes is biased towards certain taxa—and the patterns exhibited by these species might not apply to others (Caro 2001). Also, genetic diversity, has not generally been considered though may be important in terms of traits valuable to humans and for conservation (Ashley et al. 2006). Conservation in human-modified landscapes may be particularly important in conserving genetic diversity because the traditional fortress PA model may encompass relatively little, especially for plants (Atta-Krah et al. 2004).

Many authors lament erosion of ecological knowledge to maintain species, especially trees, medicinal plants, and wild food plants, and urge more effort towards domestication, cultivation, and marketing to provide farmers with the means to conserve species while easing pressure on wild stock and improving food security and economic stability (Kindt et al. 2008; Ntupanyama et al. 2008; Tabuti et al. 2009; Leakey and Tchoundjeu 2001; Dold and Cocks 2002; Dovie et al. 2007; Khumalo et al. 2012). However, care must be taken to ensure that genetic diversity is maintained in the process (Lengkeek et al. 2006;

Muchugi et al. 2008). Development of domestication and cultivation methods could promote the use of native species in human-dominated lands, and these native plants may contribute to conservation of other taxa (Dovie et al. 2007), but more research is clearly required.

### Implementing policies

Given the limitations of the available science, it is difficult to develop strategies to encourage land uses that are of the highest conservation value. The effect of policy on biodiversity conservation in human-modified landscapes under different land tenure systems and different settlement patterns needs more research because decisions are largely opinion driven and not evidence based (Homewood 2004; Duvall 2008). Perhaps the community-based conservation literature, which has focused heavily on implementation and policy, could lend some insight. A review of this literature stresses that better implementation results are achieved when there is quality governance, resilient local institutions with local power and accountability, consideration for local context, integration across social and ecological systems, and mutual learning involving communities and other involved parties, e.g. outside experts (Balint and Mashinya 2008). NGO's and foreign aid are more likely to encourage successful conservation when projects are flexible, small-scale, and targeted at local interests, and when they prioritize innovation, learning, and experimentation (Nelson 2009). Conservationists must also take cognizance of perspectives and needs of local communities in both rural and urban settings in order to better engage them in conservation management (Ferketic et al. 2010). CBC projects that are independent of PAs are excellent opportunities to maintain biodiversity on human-modified land of marginal use for agriculture; and expert opinion, monitoring, and ecological modeling tools can help communities manage their natural resources (Du Toit 2002).

We have indicated several gaps in the literature on biodiversity in African human-modified landscapes, and while much more work is required to create sensible policies that meet conservation needs and those of governments and people (Ashley et al. 2006), as it stands, current research can go some way towards supporting policy-making. Studies of biodiversity persistence in different land-use options for a given region can be incorporated into scenario modeling for future development. For example, Turpie et al. (2007) amalgamated studies of plants, invertebrates, birds, and mammals in human-modified landscapes to predict how varying levels of afforestation or dairy production in the Drakensberg grasslands of South Africa would influence alpha diversity.

Some generalities emerge from the literature that may be helpful in working towards sensible policies. Generally, diversifying human-modified landscapes at all levels, e.g. polyculture cropping, diverse agroforestry, and maintaining farmlands with high heterogeneity in terms of both crops and vegetation structure, is likely to support more species than do more homogenous land uses, while potentially also providing economic stability, given fluctuating markets for specific crops (Franzen and Mulder 2007). Endemic and specialist species tend not to persist in human-modified landscapes; thus, PAs continue to be crucial to conservation efforts, and expanding PA coverage within areas rich in such species is important (Jenkins et al. 2013). Furthermore, a regional approach to management of these sensitive species is necessary. Past and present implementation strategies are beyond the scope of this review, yet there is literature dealing with such strategies in Africa that may be of use, e.g. certification of sustainable and biodiversity friendly products (Liliehalm and Weatherly 2010).

## Living with nature

Maintaining biodiversity in landscapes where humans live, work, and extract resources implies that humans will have to coexist with other species. While the consequences of living without nature may be worse than the difficulties of living with it, certain issues present considerable, though not unmanageable, obstacles for promoting conservation beyond PAs. Many, and perhaps most, species are easily compatible with human livelihoods and may have a positive impact via aesthetic or functional value, but some, especially large mammals, can be problematic in human-dominated landscapes, e.g. carnivores threaten livelihoods by preying on livestock and, occasionally, people. Nonetheless, specific and practical actions can greatly reduce the probability of carnivore attacks. For example, in Kenyan communal lands, having a domestic dog accompany herds reduces risk of a carnivore attack by 63 %; conversely each additional boma gate increases risk of attack by 40 % (Woodroffe et al. 2007). However, carnivores are not the only concern. Other animals, such as baboons and bush pigs, can damage structures and destroy crops while larger herbivores, such as elephants, also threaten human lives. Knowledge of attitudes of people employing different land uses can help land-use planners develop strategies to reduce conflict and negative attitudes towards conservation. For example, crop agriculture should not be encouraged in predominantly pastoral areas where elephants and people coexist relatively peacefully (Gadd 2005). Furthermore, land-use planning that incorporates knowledge of which crops are most likely to generate conflict could allow creation of buffer zones in areas with high conflict potential (Hockings and McLennan 2012).

The risk of disease transmission poses an additional difficulty. Diseases of domestic animals threaten wildlife. For example, domestic dogs are carriers of canid diseases transmissible to wild carnivores (Butler et al. 2004) and were partly responsible for extinction of the African wild dog *Lycaon pictus* and decimation of lions *Panthera leo* in areas of the Serengeti (see Woodroffe 1999). Additionally, livestock can transmit animal diseases (e.g. bovine tuberculosis) to wildlife with negative conservation outcomes, while wildlife can also transmit diseases (e.g. foot and mouth) to livestock with immense economic consequences (Michel et al. 2006; Thomson 2009).

Fencing has been heavily used in Africa to assist people in their ability to coexist with nature—to reduce direct conflict and disease transmission. Laws regarding fencing differ by country; for example, Zambia requires game fences while Namibia encourages large-scale cooperation between game-farmers to discourage fencing (McGranahan 2008). Obviously, fencing has serious ecological consequences (Trimble and van Aarde 2010; Hayward and Kerley 2009) and is anathema in many ways to the goals of conservation, especially conservation beyond PAs (Trimble and van Aarde 2010; Woodroffe et al. 2014). However, non-traditional fencing technologies (see Hayward and Kerley 2009), such as fences targeted at particular problem species (e.g. elephant fences that allow other species to pass), virtual barriers, or fencing wildlife out of villages and fields instead of into PAs, may be acceptable compromises. The effect of fences on the persistence of species in human-modified landscapes certainly deserves more investigation.

Economically, wild animals provide an important resource for many people in Africa (Bharucha and Pretty 2010), which may threaten species persistence. ‘Sustainable use’ is frequently discussed with relation to bushmeat hunting, but food scarcity and population growth dictate that it will likely be impossible to enforce rules for sustainable use unless food security issues are addressed (Fa et al. 2003). Sustainable harvesting is also an issue for plants (Sambou et al. 2002). Community forests must be carefully managed, e.g. by restricting harvesting of pole-sized stems to certain species, to ensure that species are not

used to extinction (Obiri et al. 2002). Additionally, rules must be assessed to ensure that they achieve the desired goals; for example, in the Republic of Guinea, tax to the forestry administration for harvesting palm wine counterproductively encourages harvesters to employ lethal yet profitable methods of harvesting to compensate for the initial investment (Sambou et al. 2002).

## Conclusion

There is clearly both necessity and great potential for human-modified land in sub-Saharan Africa to contribute to the conservation of the continent's biodiversity. While PAs will remain essential, and are especially important for protecting species sensitive to human disturbance (Devineau et al. 2009), a greater focus on biodiversity conservation beyond their boundaries could be complementary to overall conservation goals. The information gleaned from studies of biodiversity in human-modified landscapes in Africa discussed in this review goes some way toward providing policy-makers with evidence to support defensible decisions for land-use planning and conservation management beyond PAs (see Table 1). Improving the amenity of human-modified landscapes for biodiversity can be encouraged at all levels from individuals' choices to plant indigenous home gardens, to grass roots endeavors to manage communal resources, to communities deciding to share their land with wildlife, to commercial farms going organic and maintaining patches of natural habitat. Governmental intervention at the level of the city (e.g. green space planning), region (e.g. extension agencies demonstrating biodiversity friendly agricultural practices), nation (e.g. policy-setting for control of invasive species, pesticide or poison usage, and land-use zoning), or even internationally (e.g. cooperative removal of boundary fences) are also warranted.

Although several factors including lack of knowledge, implementation challenges, and problems of coexistence with wildlife may constrain successful implementation of biodiversity conservation in human-modified landscapes, given each constraint, opportunity exists for progress. On the bright side, scientific interest in the topic is increasing (Trimble and van Aarde 2012), and as research accumulates, it will allow for systematic reviews useful for policy decisions. Additionally, many issues associated with human-wildlife coexistence are primarily related to large mammals and efforts to solve these problems should continue. Meanwhile, the barriers to implementing strategies to conserve other species groups in human-modified landscapes are far from insurmountable and such strategies should be prioritized.

**Acknowledgments** M. J. T. was supported by a National Science Foundation Graduate Research Fellowship and R. J. v. A. through various grants to the Chair in Conservation Ecology at CERU.

## References

- Abate T, van Huis A, Ampofo JKO (2000) Pest management strategies in traditional agriculture: an African perspective. *Annu Rev Entomol* 45:631–659
- Adams WM, Hulme D (2001) If community conservation is the answer in Africa, what is the question? *Oryx* 35:193–200
- Alston KP, Richardson DM (2006) The roles of habitat features, disturbance, and distance from putative source populations in structuring alien plant invasions at the urban/wildland interface on the Cape Peninsula, South Africa. *Biol Conserv* 132:183–198

- Anderson J, Rowcliffe J, Cowlshaw G (2007) Does the matrix matter? A forest primate in a complex agricultural landscape. *Biol Conserv* 135:212–222
- Armstrong AJ (2008) Translocation of black-headed dwarf chameleons *Bradypodion melanocephalum* in Durban, KwaZulu-Natal, South Africa. *Afr J Herpetol* 57:29–41
- Ashley R, Russell D, Swallow B (2006) The policy terrain in protected area landscapes: challenges for agroforestry in integrated landscape conservation. *Biodivers Conserv* 15:663–689
- Atta-Krah K, Kindt R, Skilton JN, Amaral W (2004) Managing biological and genetic diversity in tropical agroforestry. *Agrofor Syst* 61:183–194
- Augusseau X, Nikiéma P, Torquebiau E (2006) Tree biodiversity, land dynamics and farmers' strategies on the agricultural frontier of southwestern Burkina Faso. *Biodivers Conserv* 15:613–630
- Baiphethi MN, Jacobs PT (2009) The contribution of subsistence farming to food security in South Africa. *Agrekon* 48:459–482
- Balint PJ, Mashinya J (2008) CAMPFIRE during Zimbabwe's national crisis: local impacts and broader implications for community-based wildlife management. *Soc Nat Resour* 21:783–796
- Balmford A, Moore JL, Brooks T, Burgess N, Hansen LA, Williams P, Rahbek C (2001) Conservation conflicts across Africa. *Science* 291:2616–2619
- Baudron F, Corbeels M, Andersson JA, Sibanda M, Giller KE (2011) Delineating the drivers of waning wildlife habitat: the predominance of cotton farming on the fringe of protected areas in the Mid-Zambezi Valley, Zimbabwe. *Biol Conserv* 144:1481–1493
- Beresford AE, Buchanan GM, Donald PF, Butchart SHM, Fishpool LDC, Rondinini C (2011) Poor overlap between the distribution of protected areas and globally threatened birds in Africa. *Anim Conserv* 14:99–107
- Bhagwat S, Willis K, Birks H, Whittaker R (2008) Agroforestry: a refuge for tropical biodiversity? *Trends Ecol Evol* 23:261–267
- Bharucha Z, Pretty J (2010) The roles and values of wild foods in agricultural systems. *Philos Trans R Soc Lond B* 365:2913–2926
- Bigirimana J, Bogaert J, De Canniere C, Bigendako M-J, Parmentier I (2012) Domestic garden plant diversity in Bujumbura, Burundi: role of the socio-economical status of the neighborhood and alien species invasion risk. *Landsc Urban Plan* 107:118–126
- Bond WJ, Parr CL (2010) Beyond the forest edge: ecology, diversity and conservation of the grassy biomes. *Biol Conserv* 143:2395–2404
- Bothma J (ed) (2002) Game ranch management, 4th edn. Van Schaik, Pretoria
- Brink AB, Eva HD (2009) Monitoring 25 years of land cover change dynamics in Africa: a sample based remote sensing approach. *Appl Geogr* 29:501–512
- Burgess ND, Balmford A, Cordeiro NJ, Fjeldsa J, Kuper W, Rahbek C, Sanderson EW, Scharlemann JPW, Sommer JH, Williams PH (2007) Correlations among species distributions, human density and human infrastructure across the high biodiversity tropical mountains of Africa. *Biol Conserv* 134:164–177
- Butler JRA, du Toit JT, Bingham J (2004) Free-ranging domestic dogs (*Canis familiaris*) as predators and prey in rural Zimbabwe: threats of competition and disease to large wild carnivores. *Biol Conserv* 115:369–378
- Caro TM (1999) Densities of mammals in partially protected areas: the Katavi ecosystem of western Tanzania. *J Appl Ecol* 36:205–217
- Caro TM (2001) Species richness and abundance of small mammals inside and outside an African national park. *Biol Conserv* 98:251–257
- Carruthers J (2009) National Parks in South Africa. In: Suich H, Child B, Spenceley A (eds) *Evolution and innovation in wildlife conservation: parks and game ranches to transfrontier conservation areas*. Earthscan, London
- Carvalho LG, Seymour CL, Veldtman R, Nicolson SW (2010) Pollination services decline with distance from natural habitat even in biodiversity-rich areas. *J Appl Ecol* 47:810–820
- Chown SL, van Rensburg BJ, Gaston KJ, Rodrigues ASL, van Jaarsveld AS (2003) Energy, species richness, and human population size: conservation implications at a national scale. *Ecol Appl* 13:1233–1241
- Cilliers SS, Williams NSG, Barnard FJ (2008) Patterns of exotic plant invasions in fragmented urban and rural grasslands across continents. *Landsc Ecol* 23:1243–1256
- Ciofolo I (1995) West Africa's last giraffes: the conflict between development and conservation. *J Trop Ecol* 11:577–588
- Cousins JA, Sadler JP, Evans J (2008) Exploring the role of private wildlife ranching as a conservation tool in South Africa: stakeholder perspectives. *Ecol Soc* 13:Art. 43
- Cowling RM, Pressey RL (2003) Introduction to systematic conservation planning in the Cape Floristic Region. *Biol Conserv* 112:1–13

- Cox RL, Underwood EC (2011) The importance of conserving biodiversity outside of protected areas in Mediterranean ecosystems. *PLoS ONE* 6:e14508
- Craigie ID, Baillie JEM, Balmford A, Carbone C, Collen B, Green RE, Hutton JM (2010) Large mammal population declines in Africa's protected areas. *Biol Conserv* 143:2221–2228
- Cumming MS, Wesolowska W (2004) Habitat separation in a species-rich assemblage of jumping spiders (Araneae: Salticidae) in a suburban study site in Zimbabwe. *J Zool* 262:1–10
- Curtis O, Simmons RE, Jenkins AR (2004) Black Harrier *Circus maurus* of the Fynbos biome, South Africa: a threatened specialist or an adaptable survivor? *Bird Conserv Int* 14:233–245
- Daily GC, Ehrlich PR, Sanchez-Azofeifa GA (2001) Countryside biogeography: use of human-dominated habitats by the avifauna of southern Costa Rica. *Ecol Appl* 11:1–13
- Darkoh MBK (2003) Regional perspectives on agriculture and biodiversity in the drylands of Africa. *J Arid Environ* 54:261–279
- de Oliveira JAP, Balaban O, Doll CNH, Moreno-Peñaranda R, Gasparatos A, Iossifova D, Suwa A (2011) Cities and biodiversity: perspectives and governance challenges for implementing the convention on biological diversity (CBD) at the city level. *Biol Conserv* 144:1302–1313
- Dean WRJ, Milton SJ, Jeltsch F (1999) Large trees, fertile islands, and birds in arid savanna. *J Arid Environ* 41:61–78
- Deininger K, Byerlee D, Lindsay J, Norton A, Selod H, Stickler M (2011) Rising global interest in farmland: can it yield sustainable and equitable benefits?. The World Bank, Washington, DC
- Devineau JL, Fournier A, Nignan S (2009) “Ordinary biodiversity” in western Burkina Faso (West Africa): what vegetation do the state forests conserve? *Biodivers Conserv* 18:2075–2099
- UN Population Division (2012) World urbanization prospects: the 2011 revision. United Nations, Department of Economic and Social Affairs, Population Division
- Djossa BA, Fahr J, Wiegand T, Ayihouenou BE, Kalko EK, Sinsin BA (2008) Land use impact on *Vitellaria paradoxa* C.F. Gaerten. stand structure and distribution patterns: a comparison of Biosphere Reserve of Pendjari in Atacora district in Benin. *Agrofor Syst* 72:205–220
- Doak DF, Bakker VJ, Goldstein BE, Hale B (2013) What is the future of conservation? *Trends Ecol Evol* 29:77–81
- Dold AP, Cocks ML (2002) The trade in medicinal plants in the Eastern Cape Province, South Africa. *S Afr J Sci* 98:589–597
- Donald PF, Buchanan GM, Collar NJ, Abebe YD, Gabremichael MN, Mwangi MAK, Ndong'ang'a PK, Spottiswoode CN, Wondafrash M (2010) Rapid declines in habitat quality and population size of the Liben (Sidamo) Lark *Heteromirafra sidamoensis* necessitate immediate conservation action. *Bird Conserv Int* 20:1–12
- Dovie DBK, Shackleton CM, Witkowski ETF (2007) Conceptualizing the human use of wild edible herbs for conservation in South African communal areas. *J Environ Manag* 84:146–156
- Du Toit JT (2002) Wildlife harvesting guidelines for community-based wildlife management: a southern African perspective. *Biodivers Conserv* 11:1403–1416
- Dures SG, Cumming GS (2010) The confounding influence of homogenising invasive species in a globally endangered and largely urban biome: does habitat quality dominate avian biodiversity? *Biol Conserv* 143:768–777
- Duvall CS (2008) Human settlement ecology and chimpanzee habitat selection in Mali. *Landsc Ecol* 23:699–716
- Eardley CD, Gikungu M, Schwarz MP (2009) Bee conservation in Sub-Saharan Africa and Madagascar: diversity, status and threats. *Apidologie* 40:355–366
- Fa JE, Currie D, Meeuwig J (2003) Bushmeat and food security in the Congo Basin: linkages between wildlife and people's future. *Environ Conserv* 30:71–78
- Fabricius C, Burger M, Hockey PAR (2003) Comparing biodiversity between protected areas and adjacent rangeland in xeric succulent thicket, South Africa: arthropods and reptiles. *J Appl Ecol* 40:392–403
- Fairbanks DHK (2004) Regional land-use impacts affecting avian richness patterns in Southern Africa—insights from historical avian atlas data. *Agric Ecosyst Environ* 101:269–288
- Fandohan AB, Assogbadjo AE, Kakai RLG, Sinsin B, Van Damme P (2010) Impact of habitat type on the conservation status of tamarind (*Tamarindus indica* L.) populations in the W National Park of Benin. *Fruits* 65:11–19
- Ferketic JS, Latimer AM, Silander JA (2010) Conservation justice in metropolitan Cape Town: a study at the Macassar Dunes Conservation Area. *Biol Conserv* 143:1168–1174
- Fjeldsa J, Burgess ND (2008) The coincidence of biodiversity patterns and human settlement in Africa. *Afr J Ecol* 46:33–42
- Fjeldsa J, Burgess ND, Blyth S, de Klerk HM (2004) Where are the major gaps in the reserve network for Africa's mammals? *Oryx* 38:17–25

- Fox SJC, Hockey PAR (2007) Impacts of a South African coastal golf estate on shrubland bird communities. *S Afr J Sci* 103:27–34
- Franzen M, Mulder MB (2007) Ecological, economic and social perspectives on cocoa production world-wide. *Biodivers Conserv* 16:3835–3849
- Gadd ME (2005) Conservation outside of parks: attitudes of local people in Laikipia, Kenya. *Environ Conserv* 32:50–63
- Gaigher R, Samways M (2010) Surface-active arthropods in organic vineyards, integrated vineyards and natural habitat in the Cape Floristic Region. *J Insect Conserv* 14:595–605
- Gaoue OG, Ticktin T (2008) Impacts of bark and foliage harvest on *Khaya senegalensis* (Meliaceae) reproductive performance in Benin. *J Appl Ecol* 45:34–40
- Gardiner AJ, Reid RS, Kiema S (2005) Impact of land-use on butterflies in southwestern Burkina Faso. *Afr Entomol* 13:201–212
- Gardner TA, Caro T, Fitzherbert EB, Banda T, Lalbhai P (2007) Conservation value of multiple-use areas in East Africa. *Conserv Biol* 21:1516–1525
- Gardner TA, Barlow J, Sodhi NS, Peres CA (2010) A multi-region assessment of tropical forest biodiversity in a human-modified world. *Biol Conserv* 143:2293–2300
- Garrity DP, Akinnifesi FK, Ajayi OC, Weldesemayat SG, Mowo JG, Kalinganire A, Larwanou M, Bayala J (2010) Evergreen agriculture: a robust approach to sustainable food security in Africa. *Food Secur* 2:197–214
- Giliomee JH (2006) Conserving and increasing biodiversity in the large-scale, intensive farming systems of the Western Cape, South Africa. *S Afr J Sci* 102:375–378
- Gove AD, Hylander K, Nemomisa S, Shimelis A (2008) Ethiopian coffee cultivation-Implications for bird conservation and environmental certification. *Conserv Lett* 1:208–216
- Gregory NC, Sensenig RL, Wilcove DS (2010) Effects of controlled fire and livestock grazing on bird communities in East African savannas. *Conserv Biol* 24:1606–1616
- Grey-Ross R, Downs CT, Kirkman K (2009a) Reintroduction failure of captive-bred oribi (*Ourebia ourebi*). *S Afr J Wildl Res* 39:34–38
- Grey-Ross R, Downs CT, Kirkman K (2009b) Using housing estates as conservation tools: a case study in KwaZulu-Natal, South Africa. *Appl Geogr* 29:371–376
- Hackel JD (1999) Community conservation and the future of Africa's wildlife. *Conserv Biol* 13:726–734
- Hagen M, Kraemer M (2010) Agricultural surroundings support flower-visitor networks in an Afrotropical rain forest. *Biol Conserv* 143:1654–1663
- Hall JM, Gillespie TW, Mwangoka M (2011) Comparison of agroforests and protected forests in the East Usambara mountains, Tanzania. *Environ Manag* 48:237–247
- Hansen AJ, DeFries R (2007) Ecological mechanisms linking protected areas to surrounding lands. *Ecol Appl* 17:974–988
- Hayward M, Kerley G (2009) Fencing for conservation: restriction of evolutionary potential or a riposte to threatening processes? *Biol Conserv* 142:1–13
- Hemp A (2006) The banana forests of Kilimanjaro: biodiversity and conservation of the Chagga home-gardens. *Biodivers Conserv* 15:1193–1217
- Hockings KJ, McLennan MR (2012) From forest to farm: systematic review of cultivar feeding by chimpanzees-management implications for wildlife in anthropogenic landscapes. *PLoS One* 7:e33391–e33391
- Holdo RM, Galvin KA, Knapp E, Polasky S, Hilborn R, Holt RD (2010) Responses to alternative rainfall regimes and antipoaching in a migratory system. *Ecol Appl* 20:381–397
- Hole DG, Huntley B, Arinaitwe J, Butchart SHM, Collingham YC, Fishpool LDC, Pain DJ, Willis SG (2011) Toward a management framework for networks of protected areas in the face of climate change. *Conserv Biol* 25:305–315
- Homewood KM (2004) Policy, environment and development in African rangelands. *Environ Sci Policy* 7:125–143
- Homewood K, Lambin EF, Coast E, Kariuki A, Kikula I, Kivelia J, Said M, Serneels S, Thompson M (2001) Long-term changes in Serengeti-Mara wildebeest and land cover: pastoralism, population, or policies? *Proc Natl Acad Sci USA* 98:12544–12549
- IMF (2013) World economic outlook, April 2013: Hopes, realities, risks. International Monetary Fund, Washington, DC
- Jacobs CT, Scholtz CH, Escobar F, Davis ALV (2010) How might intensification of farming influence dung beetle diversity (Coleoptera: Scarabaeidae) in Maputo Special Reserve (Mozambique)? *J Insect Conserv* 14:389–399
- Jenkins C, Joppa LN (2009) Expansion of the global terrestrial protected area system. *Biol Conserv* 142:2166–2174

- Jenkins C, Pimm S, Joppa L (2013) Global patterns of terrestrial vertebrate diversity and conservation. *Proc Natl Acad Sci USA* 110:E2602–E2610
- Johnson SD, Neal PR, Peter CI, Edwards TJ (2004) Fruiting failure and limited recruitment in remnant populations of the hawkmoth-pollinated tree *Oxyanthus pyriformis* subsp *pyriformis* (Rubiaceae). *Biol Conserv* 120:31–39
- Kalaba KF, Chirwa P, Syampungani S, Ajayi OC (2010) Contribution of Agroforestry to biodiversity and livelihoods improvement in rural communities of Southern African regions. In: Tscharrntke T, Leuschner C, Veldkamp E, Faust H, Guhardja E, Bidin A (eds) *Tropical rainforests and agroforests under global change: ecological and socio-economic valuations*. Springer, Berlin
- Kareiva P, Marvier M (2012) What is conservation science? *Bioscience* 62:962–969
- Kehinde T, Samways MJ (2012) Endemic pollinator response to organic vs. conventional farming and landscape context in the Cape Floristic Region biodiversity hotspot. *Agric Ecosyst Environ* 146:162–167
- Kerley GIH, Knight MH, DeKock M (1995) Desertification of subtropical thicket in the Eastern Cape, South Africa: are there alternatives? *Environ Monit Assess* 37:211–230
- Khumalo S, Chirwa PW, Moyo BH, Syampungani S (2012) The status of agrobiodiversity management and conservation in major agroecosystems of Southern Africa. *Agric Ecosyst Environ* 157:17–23
- Kindt R, Kalinganire A, Larwanou M, Belem M, Dakouo JM, Bayala J, Kaire M (2008) Species accumulation within land use and tree diameter categories in Burkina Faso, Mali, Niger and Senegal. *Biodivers Conserv* 17:1883–1905
- Kinnaird MF, O'Brien TG (2012) Effects of private-land use, livestock management, and human tolerance on diversity, distribution, and abundance of large African mammals. *Conserv Biol* 26:1026–1039
- Kirby JS, Stattersfield AJ, Butchart SHM, Evans MI, Grimmett RFA, Jones VR, O'Sullivan J, Tucker GM, Newton I (2008) Key conservation issues for migratory land- and waterbird species on the world's major flyways. *Bird Conserv Int* 18:S49–S73
- Kiringe JW, Okello MM, Ekajul SW (2007) Managers' perceptions of threats to the protected areas of Kenya: prioritization for effective management. *Oryx* 41:314–321
- Konecny A, Koubek P, Bryja J (2010) Indications of higher diversity and abundance of small rodents in human-influenced Sudanian savannah than in the Niokolo Koba National Park (Senegal). *Afr J Ecol* 48:718–726
- Laube I, Breitbach N, Bohning-Gaese K (2008) Avian diversity in a Kenyan agroecosystem: effects of habitat structure and proximity to forest. *J Ornithol* 149:181–191
- Leakey RRB, Tchoundjeu Z (2001) Diversification of tree crops: domestication of companion crops for poverty reduction and environmental services. *Exp Agric* 37:279–296
- Lengkeek AG, Mwangi AM, Agufa CAC, Ahenda JO, Dawson IK (2006) Comparing genetic diversity in agroforestry systems with natural forest: a case study of the important timber tree *Vitex fischeri* in central Kenya. *Agrofor Syst* 67:293–300
- Liliehalm RJ, Weatherly WP (2010) Kibale forest wild coffee: challenges to market-based conservation in Africa. *Conserv Biol* 24:924–930
- Lindsey PA, Romañach SS, Davies-Mostert HT (2009) The importance of conservancies for enhancing the value of game ranch land for large mammal conservation in southern Africa. *J Zool* 277:99–105
- Lipsev MK, Hockey PAR (2010) Do ecological networks in South African commercial forests benefit grassland birds? A case study of a pine plantation in KwaZulu-Natal. *Agric Ecosyst Environ* 137:133–142
- Loarie SR, Duffy PB, Hamilton H, Asner GP, Field CB, Ackerly DD (2009) The velocity of climate change. *Nature* 462:1052–1055
- Lubbe CS, Siebert SJ, Cilliers SS (2010) Political legacy of South Africa affects the plant diversity patterns of urban domestic gardens along a socio-economic gradient. *Sci Res Essays* 5:2900–2910
- Machange RW, Jenkins AR, Navarro RA (2005) Eagles as indicators of ecosystem health: is the distribution of Martial Eagle nests in the Karoo, South Africa, influenced by variations in land-use and rangeland quality? *J Arid Environ* 63:223–243
- Mangnall MJ, Crowe TM (2003) The effects of agriculture on farmland bird assemblages on the Agulhas Plain, Western Cape, South Africa. *Afr J Ecol* 41:266–276
- McCann K, Theron LJ, Morrison K (2007) Conservation priorities for the Blue Crane (*Anthropoides paradiseus*) in South Africa—the effects of habitat changes on distribution and numbers. *Ostrich* 78:205–211
- McGeoch MA, Chown SL (1997) Impact of urbanization on a gall-inhabiting Lepidoptera assemblage: the importance of reserves in urban areas. *Biodivers Conserv* 6:979–993
- McGranahan DA (2008) Managing private, commercial rangelands for agricultural production and wildlife diversity in Namibia and Zambia. *Biodivers Conserv* 17:1965–1977

- Metzger KL, Sinclair ARE, Hilborn R, Hopcraft JGC, Mduma SAR (2010) Evaluating the protection of wildlife in parks: the case of African buffalo in Serengeti. *Biodivers Conserv* 19:3431–3444
- Michel AL, Bengis RG, Keet DF, Hofmeyr M, de Klerk LM, Cross PC, Jolles AE, Cooper D, Whyte IJ, Buss P, Godfroid J (2006) Wildlife tuberculosis in South African conservation areas: implications and challenges. *Vet Microbiol* 112:91–100
- Midega CAO, Khan ZR, van den Berg J, Ogot C, Dippenaar-Schoeman AS, Pickett JA, Wadhams LJ (2008) Response of ground-dwelling arthropods to a 'push-pull' habitat management system: spiders as an indicator group. *J Appl Entomol* 132:248–254
- Molebatsi LY, Siebert SJ, Cilliers SS, Lubbe CS, Davoren E (2010) The Tswana tshimo: a homegarden system of useful plants with a particular layout and function. *Afr J Agric Res* 5:2952–2963
- Muchugi A, Muluvi GM, Kindt R, Kadu CAC, Simons AJ, Jamnadass RH (2008) Genetic structuring of important medicinal species of genus *Warburgia* as revealed by AFLP analysis. *Tree Genet Genome* 4:787–795
- Mulwa RK, Boehning-Gaese K, Schleuning M (2012) High bird species diversity in structurally heterogeneous farmland in western Kenya. *Biotropica* 44:801–809
- Munyuli MBT (2012) Micro, local, landscape and regional drivers of bee biodiversity and pollination services delivery to coffee (*Coffea canephora*) in Uganda. *Int J Biodivers Sci Eco Serv Manag* 8:190–203
- Muriuki GW, Njoka TJ, Reid RS, Nyariki DM (2005) Tsetse control and land-use change in Lambwe valley, south-western Kenya. *Agric Ecosyst Environ* 106:99–107
- Muriuki G, McAlpine C, Seabrook L, Baxter G (2011) The role of squatters in retention of native vegetation: a case study of the Chyulu Hills, Kenya. *Appl Geogr* 31:577–589
- Myers N, Mittermeier RA, Mittermeier CG, da Fonseca GAB, Kent J (2000) Biodiversity hotspots for conservation priorities. *Nature* 403:853–858
- Mysterud A (2010) Still walking on the wild side? Management actions as steps towards 'semi-domestication' of hunted ungulates. *J Appl Ecol* 47:920–925
- NAMC (2006) Report on the investigation to identify problems for sustainable growth and development in South African Wildlife ranching. National Agricultural Marketing Council, Pretoria, South Africa
- Negash M, Yirdaw E, Luukkanen O (2012) Potential of indigenous multistrata agroforests for maintaining native floristic diversity in the south-eastern Rift Valley escarpment, Ethiopia. *Agrofor Syst* 85:9–28
- Nelson F (2009) Conservation and aid: designing more effective investments in natural resource governance reform. *Conserv Biol* 23:1102–1108
- Neuschulz EL, Botzat A, Farwig N (2011) Effects of forest modification on bird community composition and seed removal in a heterogeneous landscape in South Africa. *Oikos* 120:1371–1379
- Newmark WD (2008) Isolation of African protected areas. *Front Ecol Environ* 6:321–328
- Norris K, Asase A, Collen B, Gockowksi J, Mason J, Phalan B, Wade A (2010) Biodiversity in a forest-agriculture mosaic—the changing face of West African rainforests. *Biol Conserv* 143:2341–2350
- Ntupanyama YM, Mwase WF, Stejke B, Kwapata MB, Bokosi JM, Hvostlef-Eide AK (2008) Indigenous knowledge of rural communities in Malawi on socio-economic use, propagation, biology, biodiversity and ecology of *Uapaca kirkiana* Muell. Arg. *Afr J Biotechnol* 7:2386–2396
- Obiri J, Lawes M, Mukolwe M (2002) The dynamics and sustainable use of high-value tree species of the coastal Pondoland forests of the Eastern Cape Province, South Africa. *For Ecol Manag* 166:131–148
- O'Connor TG, Kuyler P (2009) Impact of land use on the biodiversity integrity of the moist sub-biome of the grassland biome, South Africa. *J Environ Manag* 90:384–395
- O'Farrell PJ, Milton SJ (2006) Road verge and rangeland plant communities in the southern Karoo: exploring what influences diversity, dominance and cover. *Biodivers Conserv* 15:921–938
- Olson DM, Dinerstein E, Wikramanayake ED, Burgess ND, Powell GVN, Underwood EC, D'Amico JA, Itoua I, Strand HE, Morrison JC, Loucks CJ, Allnutt TF, Ricketts TH, Kura Y, Lamoreux JF, Wetzel WW, Hedao P, Kassem KR (2001) Terrestrial ecoregions of the world: a new map of life on Earth. *Bioscience* 51:933–938
- Ouinavi C, Sokpon N (2008) Traditional agroforestry systems as tools for conservation of genetic resources of *Milicia excelsa* Welw. C.C. Berg in Benin. *Agrofor Syst* 74:17–26
- Pare S, Tigabu M, Savadogo P, Oden PC, Ouadba JM (2010) Does designation of protected areas ensure conservation of tree diversity in the Sudanian dry forest of Burkina Faso? *Afr J Ecol* 48:347–360
- Pauw A (2007) Collapse of a pollination web in small conservation areas. *Ecology* 88:1759–1769
- Pettorelli N, Lobora AL, Msuha MJ, Foley C, Durant SM (2010) Carnivore biodiversity in Tanzania: revealing the distribution patterns of secretive mammals using camera traps. *Anim Conserv* 13:131–139
- Pote J, Shackleton C, Cocks M, Lubke R (2006) Fuelwood harvesting and selection in Valley Thicket, South Africa. *J Arid Environ* 67:270–287

- Pryke SR, Samways MJ (2003) Quality of remnant indigenous grassland linkages for adult butterflies (Lepidoptera) in an afforested African landscape. *Biodivers Conserv* 12:1985–2004
- Pryke JS, Samways MJ (2009) Recovery of invertebrate diversity in a rehabilitated city landscape mosaic in the heart of a biodiversity hotspot. *Landsc Urban Plan* 93:54–62
- Pryke JS, Samways MJ (2010) Significant variables for the conservation of mountain invertebrates. *J Insect Conserv* 14:247–256
- Pye-Smith C (2010a) Fodder for a better future: how agroforestry is helping transform the lives of small-holder dairy farmers in east Africa. ICRAF Trees for Change no. 6. World Agroforestry Centre, Nairobi
- Pye-Smith C (2010b) A rural revival in tanzania: how agroforestry is helping farmers to restore woodlands in Shinyanga region. ICRAF Trees for Change no. 7. World Agroforestry Centre, Nairobi
- Raebild A, Hansen HH, Dartell J, Ky JMK, Sanou L (2007) Ethnicity, land use and woody vegetation: a case study from south-western Burkina Faso. *Agrofor Syst* 70:157–167
- Ratcliffe CS, Crowe TM (2001) The effects of agriculture and the availability of edge habitat on populations of Helmeted Guineafowl *Numida meleagris* and on the diversity and composition of associated bird assemblages in KwaZulu-Natal province, South Africa. *Biodivers Conserv* 10:2109–2127
- Reid RS, Wilson CJ, Kruska RL, Mulatu W (1997) Impacts of tsetse control and land-use on vegetative structure and tree species composition in south-western Ethiopia. *J Appl Ecol* 34:731–747
- Rosenzweig ML (2003) Reconciliation ecology and the future of species diversity. *Oryx* 37:194–205
- Rouget M, Richardson DM, Cowling RM, Lloyd JW, Lombard AT (2003a) Current patterns of habitat transformation and future threats to biodiversity in terrestrial ecosystems of the Cape Floristic Region, South Africa. *Biol Conserv* 112:63–85
- Rouget MM, Richardson DM, Cowling RM (2003b) The current configuration of protected areas in the Cape Floristic Region, South Africa—reservation bias and representation of biodiversity patterns and processes. *Biol Conserv* 112:129–145
- Rutherford MC, Powrie LW, Husted LB (2012) Plant diversity consequences of a herbivore-driven biome switch from Grassland to Nama-Karoo shrub steppe in South Africa. *Appl Veg Sci* 15:14–25
- Sambou B, Goudiaby A, Ervik F, Diallo D, Camara MC (2002) Palm wine harvesting by the Bassari threatens *Borassus aethiopicum* populations in north-western Guinea. *Biodivers Conserv* 11:1149–1161
- Samways MJ, Steytler NS (1996) Dragonfly (Odonata) distribution patterns in urban and forest landscapes, and recommendations for riparian management. *Biol Conserv* 78:279–288
- Schmitt CB, Senbeta F, Denich M, Preisinger H, Boehmer HJ (2010) Wild coffee management and plant diversity in the montane rainforest of southwestern Ethiopia. *Afr J Ecol* 48:78–86
- Schreckenberg K (1999) Products of a managed landscape: non-timber forest products in the parklands of the Bassila region, Benin. *Glob Ecol Biogeogr* 8:279–289
- Schumann K, Wittig R, Thiombiano A, Becker U, Hahn K (2010) Impact of land-use type and bark- and leaf-harvesting on population structure and fruit production of the baobab tree (*Adansonia digitata* L.) in a semi-arid savanna, West Africa. *For Ecol Manag* 260:2035–2044
- Seto KC, Gueneralp B, Hutrya LR (2012) Global forecasts of urban expansion to 2030 and direct impacts on biodiversity and carbon pools. *Proc Natl Acad Sci USA* 109:16083–16088
- Siurua H (2006) Nature above people: Rolston and “fortress” conservation in the South. *Ethics Environ* 11:71–96
- Snapp SS, Blackie MJ, Gilbert RA, Bezner-Kerr R, Kanyama-Phiri GY (2010) Biodiversity can support a greener revolution in Africa. *Proc Natl Acad Sci USA* 107:20840–20845
- Soderstrom B, Kiema S, Reid RS (2003) Intensified agricultural land-use and bird conservation in Burkina Faso. *Agric Ecosyst Environ* 99:113–124
- Songorwa AN, Buhrs T, Hughey KFD (2000) Community-based wildlife management in Africa: a critical assessment of the literature. *Nat Resour J* 40:603–643
- Spottiswoode CN, Wondafrahs M, Gabremichael MN, Abebe YD, Mwangi MAK, Collar NJ, Dolman PM (2009) Rangeland degradation is poised to cause Africa’s first recorded avian extinction. *Anim Conserv* 12:249–257
- Steytler NS, Samways MJ (1995) Biotope selection by adult male dragonflies (Odonata) at an artificial lake created for insect conservation in South Africa. *Biol Conserv* 72:381–386
- Stoner C, Caro TIM, Mduma S, Mlingwa C, Sabuni G, Borner M (2007) Assessment of effectiveness of protection strategies in Tanzania based on a decade of survey data for large herbivores. *Conserv Biol* 21:635–646
- Suh AN, Samways MJ (2005) Significance of temporal changes when designing a reservoir for conservation of dragonfly diversity. *Biodivers Conserv* 14:165–178
- Tabuti JRS (2007) The uses, local perceptions and ecological status of 16 woody species of Gadumire Sub-county, Uganda. *Biodivers Conserv* 16:1901–1915

- Tabuti JRS, Ticktin T, Arinaitwe MZ, Muwanika VB (2009) Community attitudes and preferences towards woody species: implications for conservation in Nawaikoke, Uganda. *Oryx* 43:393–402
- Taylor RD, Walker BH (1978) Comparisons of vegetation use and herbivore biomass on a Rhodesian game and cattle ranch. *J Appl Ecol* 15:565–581
- Tchabi A, Coyne D, Hountondji F, Lawouin L, Wiemken A, Oehl F (2008) Arbuscular mycorrhizal fungal communities in sub-Saharan savannas of Benin, West Africa, as affected by agricultural land use intensity and ecological zone. *Mycorrhiza* 18:181–195
- Thiollay JM (2006) Large bird declines with increasing human pressure in savanna woodlands (Burkina faso). *Biodivers Conserv* 15:2085–2108
- Thirgood S, Mosser A, Tham S, Hopcraft G, Mwangomo E, Mlengeya T, Kilewo M, Fryxell J, Sinclair ARE, Borner M (2004) Can parks protect migratory ungulates? The case of the Serengeti wildebeest. *Anim Conserv* 7:113–120
- Thomson GR (2009) Currently important animal disease management issues in sub-Saharan Africa. *Onderstepoort J Vet Res* 76:129–134
- Todd SW, Hoffman MT (2009) A fence line in time demonstrates grazing-induced vegetation shifts and dynamics in the semiarid Succulent Karoo. *Ecol Appl* 19:1897–1908
- Torquebiau E, Taylor RD (2009) Natural resource management by rural citizens in developing countries: innovations still required. *Biodivers Conserv* 18:2537–2550
- Trimble M, van Aarde R (2010) Fences are more than an issue of aesthetics. *Bioscience* 60:486
- Trimble MJ, van Aarde RJ (2011) Decline of birds in a human modified coastal dune forest landscape in South Africa. *PLoS One* 6:e16176
- Trimble MJ, van Aarde RJ (2012) Geographical and taxonomic biases in research on biodiversity in human-modified landscapes. *Ecosphere* 3:Art. 119
- Trimble MJ, van Aarde RJ (2014) Amphibian and reptile communities and functional groups over a land-use gradient in a coastal tropical forest landscape of high richness and endemism. *Anim Conserv*
- Tshiguvho TE, Dean WRJ, Robertson HG (1999) Conservation value of road verges in the semi-arid Karoo, South Africa: ants (Hymenoptera: Formicidae) as bio-indicators. *Biodivers Conserv* 8:1683–1695
- Turpie JK, O'Connor T, Mills A, Robertson H (2007) The ecological and economic consequences of changing land use in the southern Drakensberg Grasslands, South Africa. *S Afr J Econ Manag S* 10:423–441
- UN-HABITAT (2010) State of African cities 2010: governance, inequalities and urban land markets. United Nations Human Settlements Programme (UN-HABITAT), Nairobi
- van Aarde R, Jackson T (2007) Megaparks for metapopulations: addressing the causes of locally high elephant numbers in southern Africa. *Biol Conserv* 134:289–297
- van Rensburg BJ, Peacock DS, Robertson MP (2009) Biotic homogenization and alien bird species along an urban gradient in South Africa. *Landsc Urban Plan* 92:233–241
- Vanderpost C (2006) Pathways of human sprawl in wilderness buffer zones. *Popul Environ* 27:285–306
- Venter SM, Witkowski ETF (2010) Baobab (*Adansonia digitata* L.) density, size-class distribution and population trends between four land-use types in northern Venda, South Africa. *For Ecol Manag* 259:294–300
- Von Hase A, Rouget M, Cowling RM (2010) Evaluating private land conservation in the Cape Lowlands, South Africa. *Conserv Biol* 24:1182–1189
- Waltert M, Bobo KS, Kaupa S, Montoya ML, Nsanyi MS, Fermon H (2011) Assessing conservation values: biodiversity and endemism in tropical land use systems. *PLoS One* 6:e16238
- Watkinson AR, Ormerod SJ (2001) Grasslands, grazing and biodiversity: editors' introduction. *J Appl Ecol* 38:233–237
- Wessels KJ, Reyers B, van Jaarsveld AS, Rutherford MC (2003) Identification of potential conflict areas between land transformation and biodiversity conservation in north-eastern South Africa. *Agric Ecosyst Environ* 95:157–178
- Western D, Groom R, Worden J (2009) The impact of subdivision and sedentarization of pastoral lands on wildlife in an African savanna ecosystem. *Biol Conserv* 142:2538–2546
- Whitmore C, Crouch TE, Slotow RH (2002) Conservation of biodiversity in urban environments: invertebrates on structurally enhanced road islands. *Afr Entomol* 10:113–126
- Wildlife Conservation Society and Center for International Earth Science Information Network (2005) Last of the Wild Project, Version 2 (LWP-2): Global human influence index (HII) dataset (geographic). NASA Socioeconomic Data and Applications Center (SEDAC). <http://sedac.ciesin.columbia.edu/data/set/wildareas-v2-human-influence-index-geographic>. Accessed 6 Nov 2013
- Witt ABR, Samways MJ (2004) Influence of agricultural land transformation and pest management practices on the arthropod diversity of a biodiversity hotspot, the Cape Floristic Region, South Africa. *Afr Entomol* 12:89–95

- Wittemyer G, Elsen P, Bean WT, Burton ACO, Brashares JS (2008) Accelerated human population growth at protected area edges. *Science* 321:123–126
- Woodroffe R (1999) Managing disease threats to wild mammals. *Anim Conserv* 2:185–193
- Woodroffe R, Frank LG, Lindsey PA, Ranah S, Romanach S (2007) Livestock husbandry as a tool for carnivore conservation in Africa's community rangelands: a case-control study. *Biodivers Conserv* 16:1245–1260
- Woodroffe R, Hedges S, Durant SM (2014) To fence or not to fence. *Science* 344:46–48